
HEALTH CRANICE.

Indian Health Service


Division of Epidemiology and Disease Prevention Influenza Surveillance Report

IHS Influenza Awareness System (IIAS) #27 (Week 14: 4/1/2012 – 4/7/2012)

Percentage of Visits for Influenza-like Illness (ILI) per week Overall ILI % (ILI visits per week / Total visits per week) (ILI) in IHS, October through April




All data are preliminary and may change as more reports are received.



% Visits for ILI Compared to Previous Week

| IHS Area | % in week 13 | % in week 14 |
|-------------|--------------|--------------|
| NATIONAL | 1.3 | 1.2 |
| Aberdeen | 1.2 | 1.4 |
| Alaska | 0.7 | 0.3 |
| Albuquerque | 1.9 | 1.8 |
| Bemidji | 1.6 | 1.1 |
| Billings | 1.5 | 1.0 |
| California | 0.8 | 0.8 |
| Nashville | 0.8 | 1.4 |
| Navajo | 3.8 | 2.8 |
| Oklahoma | 0.4 | 0.3 |
| Phoenix | 1.5 | 1.5 |
| Portland | 1.1 | 0.8 |
| Tucson | 0.4 | 0.4 |

Percentage of Visits for Influenza-like Illness (ILI) per week Overall ILI % (ILI visits per week / Total visits per week) (ILI) in IHS, 03/23/2009 through 04/07/2012




Year - Week

IHS Influenza Vaccination Surveillance

 As of 4/7/2012, sites reporting to the IIAS have administered 401,195 doses (1st and 2nd) of seasonal influenza vaccine to Active Clinical* patients.

Influenza Vaccine Coverage, IHS Influenza Awareness System Weeks 39 (09/25/2011) through 14 (04/07/2012), 2011 - 2012



(Active Clinical Population covered by the IIAS) (*Includes all doses administered.

Influenza Vaccination Coverage by Age Group

| Age Group | Active Clinical Population | 1 dose Seasonal Flu Vaccine | % Seasonal flu (1 dose) | 2 dose Seasonal Flu Vaccine | % Seasonal flu (2 dose) |
|-------------------------------|-------------------------------|-----------------------------------|-------------------------------|-----------------------------------|-------------------------------|
| 6-23 months | 27,701 | 18,998 | 68.6 | 7,248 | 26.2 |
| 2-4 years | 72,369 | 26,070 | 36.0 | 3,168 | 4.4 |
| 5-17 years | 247,084 | 81,822 | 33.1 | | |
| 18-49 years | 535,434 | 139,187 | 26.0 | | |
| 50-64 years | 196,663 | 76,713 | 39.0 | | • |
| 65+ years | 108,686 | 47,989 | 44.2 | | |
| Children (6 mnths - 17 years) | 347,154 | 126,890 | 36.6 | | |
| Adults (18+) | 840,783 | 263,889 | 31.4 | | • |
| All(6 mnths +) | 1,187,937 | 390,779 | 32.9 | • | • |

^{*} Active Clinical population is defined as patients who have had at least 2 visits in the last 3 years

** Limited to sites that exported both doses administered and active clinical population data

Influenza Vaccination Coverage by Risk Group

| Risk Group | Active Clinical Population | 1 dose Seasonal Flu Vaccine | % Seasonal flu (1 dose) |
|---------------|-------------------------------|-----------------------------------|-------------------------------|
| Asthma | 137,448 | 23,414 | 17.0 |
| Diabetes | 157,994 | 48,746 | 30.9 |
| Pregnant | 9,006 | 3,305 | 36.7 |

^{*} Active Clinical population is defined as patients who have had at least 2 visits in the last 3 years

** Limited to sites that exported both doses administered and active clinical population data

Summary

• This report includes data from the following # of RPMS databases:

| IHS Area | # of RPMS Databases Reporting |
|-------------|-------------------------------|
| Aberdeen | 17 |
| Alaska | 11 |
| Albuquerque | 6 |
| Bemidji | 3 |
| Billings | 9 |
| California | 16 |
| Nashville | 14 |
| Navajo | 7 |
| Oklahoma | 19 |
| Phoenix | 18 |
| Portland | 21 |
| Tucson | 1 |
| All Areas | 142 |

• ILI activity continues to decrease nationwide for both IHS and the general U.S.