DEPARTMENT OF HEALTH & HUMAN SERVICES

SO CHANGE WAS A

Indian Health Service 5300 Homestead Rd. NE Albuquerque, NM 87110

To: Director of IHS Maternal & Child Health Program
IHS Area Chief Medical Officers
IHS Area Immunization Coordinators
Tribal Epidemiology Centers

From: Amy Groom, IHS Immunization Program Manager

Subject: Fourth Quarter FY 2013 Immunization Report (July 1, 2013 – September 30, 2013)

Date: December 6, 2013

The Indian Health Service, Tribal and Urban Indian immunization programs report on the immunization status of American Indian and Alaska Native (Al/AN) children 3-27 months of age, 19 – 35 months of age, and Al/AN adolescents 13 – 17 years of age. They also report on influenza vaccine coverage for all age groups. These reports are submitted to the IHS Division of Epidemiology and Disease Prevention on a quarterly basis.

3-27 Month Old Report

The 3 – 27 month report is designed to help programs ensure timely vaccination and identify children who may be falling behind so they can be brought up to date. The criteria listed below are used to monitor coverage in the following age groups:

Age Group	Vaccines	Age Group	Vaccines
3-4 months	 DTAP1 IPV 1 Hib 1 Hepatitis B 1 PCV1 (ROTA1)[†] 	16-18 months	 DTAP 3 IPV 2 MMR 1 Hib 3 Hepatitis B 2 Varicella PCV4 (ROTA3)[†]
5-6 Months	 DTAP 2 IPV 2 Hib 2 Hepatitis B 2 PCV2 (ROTA2)[†] 	19-23 months	 DTAP 4 IPV 3 MMR 1 Hib 3 Hepatitis B 3 Varicella PCV4 (ROTA3)[†]
7-15 months	 DTAP 3 IPV 2 Hib 2 Hepatitis B 2 PCV3 (ROTA3)[†] 	24-27 months	 DTAP 4 IPV 3 MMR 1 Hib 3 Hepatitis B 3 Varicella PCV4 (ROTA3)[†] (Hepatitis A1)[†]


[†] Not included in Age appropriate immunization coverage calculations

Rotavirus vaccine (ROTA) and hepatitis A, though recommended, are not included in overall age appropriate coverage calculations. Data on ROTA and hepatitis A vaccine coverage are reported where available.

Summary

For FY 2013 Q4, 12 IHS Areas submitted immunization reports from IHS, Tribal, and Urban Indian health centers (I/T/U). This composite report provides information on the immunization status of 25,576 children 3-27 months old. Of these, 18,781 or 73% received all age-appropriate vaccinations compared to 73% (21,631/29,482) of children 3-27 months in the 3rd Quarter of FY 2013 [Figure 1]. Coverage by Area is included in the spreadsheet.

Figure 1



Two Year Old Report


The Two year old reports (children 19-35 months old) are used to monitor progress towards the GPRA childhood immunization indicator.


Starting in 2011, the vaccines series measure used to monitor coverage in this age group is, the 4:3:1:3:3:1:4 (4 DTaP, 3 IPV, 1 MMR, 3 Hib, 3 Hep B, 1 VAR, 4 PCV) series. The HP 2020 goal is 80% with the 4:3:1:3:3:1:4 series, and 90% coverage with each individual vaccine in the series.

4:3:1:3:3:1:4 Series

In FY 2013 Q4, 12 IHS Areas completed reports. Overall 74% (13,393/18,098) of two year olds had completed the 4:3:1:3:3:1:4 vaccine series compared to 74% (14,511/19,510) in FY 2013 Q3. Coverage for both series by Area is included in the spreadsheet.

Figure 2





ADOLESCENT REPORT

This report collects data on adolescents 13 – 17 years who meet the "Active Clinical User" definition (e.g. 2 visits in the last 3 years) and is designed to monitor uptake of "recently" recommended adolescent vaccines (e.g. tetanus toxoid, reduced diphtheria toxoid and acellular pertussis [Tdap], meningococcal conjugate [MCV4], and human papillomavirus [HPV] vaccines) and monitor coverage in the adolescent population with the following recommended childhood vaccines: 1 dose of Td or Tdap, 3 doses of hepatitis B, 2 doses of MMR, and 2 doses of Varicella/Hx of chickenpox. A summary of immunization coverage for 13 year olds and 13 –17 year olds with these vaccines is included below.


Immunization Coverage with Tdap and MCV4 for 13 year olds, Males and Females

For FY 2013 Q4, the 12 IHS Areas combined reported on 13,167 thirteen year olds compared to 15,041 thirteen year olds in the FY 2013 Q3 report. For FY 2013 Q4, 91% (11,933/13,167) received 1 dose of Tdap vaccine, 91% (12,027/13,167) received 1 dose of Tdap/Td vaccine, 87% (11,491/13,167) had received 1 dose of MCV4 vaccine, 79% of females received HPV 1 dose (5,222/6,617) and 63% of males received HPV 1 dose (4,139/6,612) [Figure 3].

In FY 2013 Q4, coverage by Area ranged from 84% - 96% for Tdap, 85% - 96% for Tdap/Td, and 73% - 95% for MCV4. Coverage by Area is included in the spreadsheet.

Tdap, Tdap/Td, and MCV4 Coverage for 13 year olds

Figure 3





Immunization Coverage for 13 – 17 year olds, Males and Females

For FY 2013 Q4, the 12 IHS Areas combined reported on 64,692 13–17 year olds compared to the FY 2013 Q3 report which included 73,414 13–17 year olds.

For FY 2013 Q3, 95% (61,323/64,692) received 3 doses of hepatitis B vaccine, 95% (61,244/64,692) received 2 doses of MMR vaccine, 89% (57,879/64,692) received 2 doses of Varicella vaccine or had a documented history of chickenpox, 92% (59,500/64,692) received 1 dose of Tdap vaccine, 93% (60,177/64,692) received 1 dose of Tdap or Td and 89% (57,712/64,692) received 1 dose of MCV4 vaccine. Coverage by Area is included in the spreadsheet [Figure 4].

Figure 4





HPV Coverage for 13 – 17 year olds, Females Only


For FY 2013 Q4, the 12 IHS Areas combined reported on 33,117 females 13 – 17 years of age compared to FY 2013 Q3 which included 37,131 females 13-17 years. For FY 2013 Q4, 83% (27,437/33,117) had received 1 dose of HPV, 72% (23,789/33,117) had received 2 doses of HPV and 59% (19,387/33,117) had received 3 doses of HPV [Figure 5].

For FY 2013 Q4, coverage by Area ranged from 71% - 91% for HPV1, 57% - 83% for HPV2, and 43% - 71% for HPV 3. Coverage by Area is included in the spreadsheet.

Figure 5

HPV Coverage for Females 13 - 17 years

FY10 Q4 - FY13 Q4





HPV Coverage for 13 – 17 year olds, Males Only

For FY 2013 Q4, the 12 IHS Areas combined reported on 31,934 males 13 – 17 years of age compared to FY 2013 Q3 which included 36,053 males 13-17 years. For FY 2013 Q3, 60% (19,215/31,934) had received 1 dose of HPV, 39% (12,397/31,934) had received 2 doses of HPV and 23% (7,318/31,934) had received 3 doses of HPV [Figure 6].

For FY 2013 Q4, coverage by Area ranged from 36% - 74% for HPV1, 15%-53% for HPV2, and 5% – 35% for HPV 3. Coverage by Area is included in the spreadsheet.

Figure 6





Adult Immunization Report

The Adult Immunization Report was collected for the first time FY12 Q4. This report collects data on patients meeting the "Active Clinical User "definition (e.g. 2 visits in the last 3 years) and reports vaccination rates for the following vaccines and age groups: Tdap in those 19 years+; HPV 1, 2 and 3 doses in Females 19-26 years: HPV 1, 2, and 3 doses in Males 19-21 years; Zoster vaccine in those 60 years+, and Pneumococcal polysaccharide vaccine at or after age 65 years.

For FY13 Q4 data were collected from all 12 IHS Areas on 513,079 active clinical patients 19 years and older compared to FY13 Q3 which included 564,701. Below is a chart with the coverage for each of the adult vaccines. Area specific vaccine coverage is included in the spreadsheet.

