



RESOURCE AND PATIENT MANAGEMENT SYSTEM

# **IHS Clinical Reporting System**

(BGP)

## **National GPRA/GPRAMA Report Performance Measure List and Definitions**

Version 25.0  
November 2024

Office of Information Technology (OIT)  
Division of Information Technology

# Table of Contents

|            |   |          |
|------------|---|----------|
| <b>1.0</b> | <b>CRS 2025 National GPRA/GPRAMA Report .....</b>                                       | <b>1</b> |
| 1.1        | Performance Measures .....  | 1        |
| 1.2        | CRS Denominator Definitions .....   | 6        |
| 1.2.1      | For All Denominators .....  | 6        |
| 1.2.2      | Active Clinical Population .....  | 6        |
| 1.2.3      | User Population .....   | 7        |
| 1.2.4      | Active Clinical Plus Behavioral Health (BH) Population .....                            | 7        |
| <b>2.0</b> | <b>Performance Measure Topics and Definitions .....</b>                                 | <b>9</b> |
| 2.1        | Diabetes Group .....  | 9        |
| 2.1.1      | Diabetes Prevalence .....   | 9        |
| 2.1.2      | Diabetes: Glycemic Control .....  | 10       |
| 2.1.3      | Diabetes: Blood Pressure Control .....  | 11       |
| 2.1.4      | Statin Therapy to Reduce Cardiovascular Disease Risk in Patients with<br>Diabetes ..... | 14       |
| 2.1.5      | Diabetes: Nephropathy Assessment .....  | 22       |
| 2.1.6      | Diabetic Retinopathy .....  | 24       |
| 2.2        | Dental Group .....  | 27       |
| 2.2.1      | Access to Dental Services .....   | 27       |
| 2.2.2      | Dental Sealants .....   | 28       |
| 2.2.3      | Topical Fluoride .....  | 30       |
| 2.3        | Immunization Group .....  | 31       |
| 2.3.1      | Influenza .....   | 31       |
| 2.3.2      | Adult Immunizations .....   | 33       |
| 2.3.3      | Childhood Immunizations .....   | 38       |
| 2.4        | Cancer Screening Group .....  | 50       |
| 2.4.1      | Cervical Cancer Screening .....   | 50       |
| 2.4.2      | Cancer Screening: Mammogram Rates .....   | 54       |
| 2.4.3      | Colorectal Cancer Screening .....   | 56       |
| 2.4.4      | Tobacco Use and Exposure Assessment .....   | 59       |
| 2.4.5      | Tobacco Cessation .....   | 61       |
| 2.5        | Behavioral Health Group .....   | 65       |
| 2.5.1      | Alcohol Screening .....   | 65       |
| 2.5.2      | Screening, Brief Intervention, and Referral to Treatment (SBIRT) .....                  | 68       |
| 2.5.3      | Intimate Partner (Domestic) Violence Screening .....                                    | 71       |
| 2.5.4      | Depression Screening .....  | 73       |
| 2.6        | Cardiovascular Disease Related Group .....  | 75       |
| 2.6.1      | Childhood Weight Control .....  | 75       |
| 2.6.2      | Controlling High Blood Pressure – Million Hearts .....                                  | 76       |

---

|       |   |           |
|-------|---|-----------|
| 2.6.3 | Statin Therapy for the Prevention and Treatment of Cardiovascular Disease ..... | 80        |
| 2.7   | STD-Related Group .....   | 87        |
| 2.7.1 | HIV Screening .....   | 87        |
| 2.8   | Other Clinical Group .....  | 91        |
| 2.8.1 | Breastfeeding Rates .....   | 91        |
|       | <b>Acronyms</b> .....   | <b>94</b> |
|       | <b>Contact Information</b> .....  | <b>97</b> |

## 1.0 CRS 2025 National GPRA/GPRAMA Report

### 1.1 Performance Measures

The following performance measures (Table 1-1) will be reported in the Clinical Reporting System (CRS) 2025 National Government Performance and Results Act of 1993 (GPRA)/GPRA Modernization Act (GPRAMA) Report.

Table 1-1: Notations in document

| Notation           | Location                                 | Meaning  |
|--------------------|--|--|
| GPRA:              | Preceding a measure                      | An official GPRA measure reported in the National GPRA Report submitted to Office of Management and Budget (OMB) and Congress in the annual Indian Health Service (IHS) budget process.                    |
| GPRAMA:            | Preceding a measure                      | An official GPRAMA measure reported in the National GPRA Report submitted to OMB and Congress and included in the annual Health and Human Services (HHS) Online Performance Appendix.                      |
| Plus Sign (+)      | Preceding a measure                      | The measure is <i>not</i> an official GPRA measure but <i>is included</i> in the National GPRA Report provided to OMB and Congress to provide context to a GPRA measure(s).                                |
| Section Symbol (§) | Preceding a measure                      | The measure is <i>not</i> an official GPRA measure and <i>is not included</i> in the National GPRA Report provided to OMB and Congress. Included in this document to provide context to a GPRA measure(s). |
| Asterisk (*)       | Anywhere in a code (CPT, POV, Edu, Etc.) | A “wildcard” character indicating that the code given has one or more additional characters at this location.  |
| Brackets ([ ])     | In logic definitions                     | Contains the name of the taxonomy where the associated codes reside.   |

#### DIABETES GROUP

- DIABETES PREVALENCE
  - +Diabetes Diagnosis Ever
  - §Diabetes Diagnosis during GPRA Year
- GLYCEMIC CONTROL
  - +Documented Alc

- GPRA: Poor Glycemic Control
- §A1c greater than or equal to ( $\geq$ ) 7 and less than ( $<$ ) 8
- Good Glycemic Control
- A1c less than ( $<$ ) 7
- BLOOD PRESSURE CONTROL
  - §Blood Pressure (BP) Assessed
  - GPRA: Controlled BP (less than [ $<$ ] 140/90)
- STATIN THERAPY TO REDUCE CARDIOVASCULAR DISEASE RISK IN PATIENTS WITH DIABETES
  - GPRA: Statin Therapy
  - §With Denominator Exclusions
- NEPHROPATHY ASSESSMENT
  - GPRA: Estimated Glomerular Filtration Rate (GFR) and Quantitative Urine Albumin-to-Creatinine Ratio (UACR) or History of End Stage Renal Disease (ESRD)
- RETINOPATHY ASSESSMENT
  - GPRA: Retinopathy Evaluation (No Refusals)

#### **DENTAL GROUP**

- ACCESS TO DENTAL
  - GPRA: Annual Dental Visit (No Refusals)
- DENTAL SEALANTS
  - GPRA: Dental Sealants (rate)
  - §Dental Sealants (No Refusals; count; not rate)
- TOPICAL FLUORIDE
  - GPRA: Topical Fluoride (rate)
  - §Topical Fluoride Application (No Refusals; count; not rate)

#### **IMMUNIZATIONS**

- INFLUENZA
  - GPRA: Influenza Immunization 6 months–17 years
  - GPRAMA: Influenza Immunization 18 years and older
- ADULT IMMUNIZATIONS
  - §1 Tdap/Td past 10 years
  - §1 Tdap ever

- §1 Influenza
- §2 Shingrix
- §1 up-to-date Pneumococcal
- §1:1:1 (Tdap/Td, Tdap, Influenza)
- §1:1 (Tdap/Td, Tdap)
- §1:1:1:2 (Tdap/Td, Tdap, Influenza, 2 Shingrix)
- §1:1:2 (Tdap/Td, Tdap, 2 Shingrix)
- §1:1:1:2:1 (Tdap/Td, Tdap, Influenza, 2 Shingrix, up-to-date Pneumococcal)
- §1:1:2:1 (Tdap/Td, Tdap, 2 Shingrix, up-to-date Pneumococcal)
- GPRA: All Age-appropriate Immunizations
- CHILDHOOD IMMUNIZATIONS (19 THROUGH 35 MONTHS)
  - §Active Clinical Patients with 4:3:1:3\*:3:1:4 (No Refusals)
  - GPRA: Active IMM Patients with 4:3:1:3\*:3:1:4 (No Refusals)
  - §4 DTaP
  - §3 Polio
  - §1 MMR
  - §3 or 4 HiB
  - §3 Hepatitis B
  - §1 Varicella
  - §4 Pneumococcal

### **CANCER SCREENING**

- CERVICAL CANCER SCREENING
  - GPRA: Pap smear in past three years or for ages 30 and older, Pap and HPV or HPV Primary in past five years (No Refusals)
- MAMMOGRAM RATES
  - GPRA: Mammogram (No Refusals)
- COLORECTAL CANCER SCREENING
  - GPRA: Fecal Occult Blood Test (FOBT) or Fecal Immunochemical Test (FIT) during Report Period, Flexible Sigmoidoscopy or CT colonography in past five years, Colonoscopy in past 10 years, or FIT-DNA in past three years (No Refusals)
  - §FOBT or FIT
- TOBACCO USE AND EXPOSURE ASSESSMENT
  - §Tobacco Assessment

- §Tobacco Users
  - §Smokers
  - §Smokeless Users
  - §Other Substance Users
  - §E-Cigarette Users
- §Exposed to Environmental Tobacco Smoke (ETS)
- TOBACCO CESSATION
  - §Tobacco Cessation Counseling or Smoking Cessation Aid (No Refusals)
  - §Quit Tobacco Use
  - GPRA: Tobacco Cessation Counseling, Smoking Cessation Aid, or Quit Tobacco Use

## **BEHAVIORAL HEALTH**

- ALCOHOL SCREENING
  - GPRA: Alcohol Screening 9–75 (No Refusals)
  - §Alcohol-Related Education
  - §Positive Alcohol Screen
- SCREENING, BRIEF INTERVENTION, AND REFERRAL TO TREATMENT
  - §Screening for Alcohol Use
  - §Positive Screen
  - GPRA: Brief Negotiated Interview/Brief Intervention (BNI/BI)
  - §Referral to Treatment
- INTIMATE PARTNER VIOLENCE/DOMESTIC VIOLENCE (IPV/DV) SCREENING
  - GPRAMA: Intimate Partner Violence/Domestic Violence (IPV/DV) Screening (No Refusals)
- DEPRESSION SCREENING
  - GPRA: Depression Screening or Mood Disorder Diagnosis age 12–17 (No Refusals)
  - GPRA: Depression Screening or Mood Disorder Diagnosis age 18 and older (No Refusals)
  - §Depression Screening
  - §Mood Disorder Diagnosis

## **CARDIOVASCULAR DISEASE-RELATED**

- CHILDHOOD WEIGHT CONTROL

- GPRA: Body Mass Index (BMI) 95% and Up
- CONTROLLING HIGH BLOOD PRESSURE – MILLION HEARTS
  - GPRA: BP less than (<) 140/90
- STATIN THERAPY FOR THE PREVENTION AND TREATMENT OF CARDIOVASCULAR DISEASE
  - GPRA: Statin Therapy
  - §With Denominator Exclusions

### **STD GROUP**

- HIV SCREENING
  - §Prenatal HIV Screening (No Refusals)
  - GPRA: HIV Screen Ever (No Refusals)

### **OTHER CLINICAL**

- BREASTFEEDING RATES
  - Patients 30 through 394 days of age screened for infant feeding choice (IFC) at least once
  - Patients 30 through 394 days of age screened for IFC at the age of 2 months
  - Patients 30 through 394 days of age screened for IFC at the age of 6 months
  - Patients 30 through 394 days of age screened for IFC at the age of 9 months
  - Patients 30 through 394 days of age screened for IFC at the age of 1 year
  - GPRA: Patients 30 through 394 days of age who were exclusively or mostly breastfed at 2 months of age
  - Patients 30 through 394 days of age who were exclusively or mostly breastfed at 6 months of age
  - Patients 30 through 394 days of age who were exclusively or mostly breastfed at 9 months of age
  - Patients 30 through 394 days of age who were exclusively or mostly breastfed at the age of 1 year

**Note:** Definitions for all performance measure topics included in CRS begin in Section 2.0. Definitions for numerators and denominators that are preceded by “GPRA” represent measures that are reported to OMB and Congress.



## 1.2 CRS Denominator Definitions

### 1.2.1 For All Denominators

- All patients with name “DEMO,PATIENT” or who are included in the RPMS Demo/Test Patient Search Template (the DPST option located in the Patient Care Component [PCC] Management Reports, Other section) will be excluded automatically for all denominators.
- For all measures, except as noted, patient age is calculated as of the beginning of the Report Period.

### 1.2.2 Active Clinical Population

#### 1.2.2.1 National GPRA/GPRAMA Reporting

- Must have two visits to medical clinics in the past three years prior to the end of the Report Period. Chart reviews and telephone calls from these clinics do not count; the visits must be face to face. At least one visit must be to a core medical clinic. Refer to the *Clinical Reporting System (CRS) for FY2025 Clinical Measures User Manual* for listing of these clinics.
- Must be alive on the last day of the Report Period.
- Must be American Indian/Alaska Native (AI/AN); defined as Beneficiary 01.
- Must reside in a community specified in the site’s GPRA community taxonomy, defined as all communities of residence in the defined Purchased and Referred Care (PRC) catchment area.

#### 1.2.2.2 Local Reports

- Must have two visits to medical clinics in the past three years prior to the end of the Report Period. Chart reviews and telephone calls from these clinics do not count; the visits must be face to face. At least one visit must be to a core medical clinic. Refer to the *Clinical Reporting System for FY2025 Clinical Measures User Manual* for listing of these clinics.
- Must be alive on the last day of the Report Period.
- User defines population type: AI/AN patients only, non-AI/AN, or both.
- User defines general population: single community, group of multiple communities (community taxonomy), user-defined list of patients (patient panel), or all patients regardless of community of residence.

## 1.2.3 User Population

### 1.2.3.1 National GPRA/GPRAMA Reporting

- Must have been seen at least once in the three years prior to the end of the Report Period, regardless of the clinic type, and the visit must be either ambulatory (including day surgery or observation), a hospitalization or a telemedicine visit; the rest of the service categories are excluded.
- Must be alive on the last day of the Report Period.
- Must be AI/AN; defined as Beneficiary 01.
- Must reside in a community specified in the site's GPRA community taxonomy, defined as all communities of residence in the defined PRC catchment area.

### 1.2.3.2 Local Reports

- Must have been seen at least once in the three years prior to the end of the Report Period, regardless of the clinic type, and the visit must be either ambulatory (including day surgery or observation) or a hospitalization; the rest of the service categories are excluded.
- Must be alive on the last day of the Report Period.
- User defines population type: AI/AN patients only, non-AI/AN, or both.
- User defines general population: single community, group of multiple communities (community taxonomy), user-defined list of patients (patient panel), or all patients regardless of community of residence.

## 1.2.4 Active Clinical Plus Behavioral Health (BH) Population

### 1.2.4.1 National GPRA/GPRAMA Reporting

- Must have two visits to medical or behavioral health clinics in the past three years prior to the end of the Report Period. Chart reviews and telephone calls from these clinics do not count; the visits must be face to face. At least one visit must be to a core medical clinic. Refer to the *Clinical Reporting System (CRS) for FY2025 Clinical Measures User Manual* for a listing of these clinics.
- Must be alive on the last day of the Report Period.
- Must be AI/AN; defined as Beneficiary 01.
- Must reside in a community specified in the site's GPRA community taxonomy, defined as all communities of residence in the defined PRC catchment area.

### 1.2.4.2 Local Reports

- Must have two visits to medical or behavioral health clinics in the past three years prior to the end of the Report Period. Chart reviews and telephone calls from these clinics do not count; the visits must be face to face. At least one visit must be to a core medical clinic. Refer to the *Clinical Reporting System (CRS) for FY2025 Clinical Measures User Manual* for a listing of these clinics.
- Must be alive on the last day of the Report Period.
- User defines population type: AI/AN patients only, non-AI/AN, or both.
- User defines general population: single community, group of multiple communities (community taxonomy), user-defined list of patients (patient panel), or all patients regardless of community of residence.

## 2.0 Performance Measure Topics and Definitions

The following sections define the performance measure topics and their definitions that are included in the *CRS 2025 version 25.0 National GPRA/GPRAMA Report*.

### 2.1 Diabetes Group

#### 2.1.1 Diabetes Prevalence

##### 2.1.1.1 Owner and Contact

Diabetes Program: Carmen Hardin

##### 2.1.1.2 National Reporting

NATIONAL (included in IHS Performance Report; *not* reported to OMB and Congress)

##### 2.1.1.3 Denominators

1. User Population patients.

##### 2.1.1.4 Numerators

1. Patients diagnosed with diabetes at any time before the end of the Report Period.
2. Patients diagnosed with diabetes during the Report Period.

##### 2.1.1.5 Definitions

###### Diabetes Diagnosis

At least one Purpose of Visit (POV) diagnosis recorded in the V POV file or Problem List Entry where the status is not Deleted:

- International Classification of Diseases (ICD)-9: 250.00 through 250.93 or ICD-10: E10.\* through E13.\* [SURVEILLANCE DIABETES]
- SNOMED data set PXR DIABETES (Problem List only)

For DM diagnosis during the Report Period, Problem List Entry must have a Date of Onset during the Report Period or, if no Date of Onset, then Date Entered during the Report Period.

##### 2.1.1.6 Patient Lists

Diabetic patients with most recent diagnosis.

## 2.1.2 Diabetes: Glycemic Control

### 2.1.2.1 Owner and Contact

Diabetes Program: Carmen Hardin

### 2.1.2.2 National Reporting

NATIONAL (included in IHS Performance Report; reported to OMB and Congress)

### 2.1.2.3 Denominators

1. GPRA: User Pop Diabetic patients, defined as User Population patients diagnosed with diabetes prior to the Report Period, *and* at least two visits during the Report Period, *and* two DM-related visits ever *or* DM entry on the Problem List. The key denominator for this and all diabetes-related topics follows.

### 2.1.2.4 Numerators

1. Hemoglobin A1c documented during the Report Period.
2. GPRA: Poor control: A1c greater ( $>$ ) than 9.
3. A1c is greater than or equal ( $\geq$ ) to 7 and less than ( $<$ ) 8
4. Good control: A1c less than ( $<$ ) 8.

### 2.1.2.5 Definition

#### Diabetes

First DM Purpose of Visit (POV) recorded in the V POV file or Problem List Entry where the status is not Deleted with Date of Onset or Date Entered prior to the Report Period:

- ICD-9: 250.00 through 250.93 or ICD-10: E10.\* through E13.\*  
[SURVEILLANCE DIABETES]
- SNOMED data set PXRDM DIABETES (Problem List only)

#### A1c

Searches for most recent A1c test with a result during the Report Period. If more than one A1c test is found on the same day or the same visit and one test has a result and the other does not, the test with the result will be used. If both tests have a result, the last test done on the visit will be used.

If an A1c test with a result is not found, CRS searches for the most recent A1c test without a result.

- A1c defined as any of the following:
  - Current Procedural Terminology (CPT) 83036, 83037, 3044F to 3046F, 3047F (old code), 3051F, 3052F [BGP HGBA1C CPTS]
  - Logical Observations Identifiers, Names, Codes (LOINC) taxonomy: 17855-8, 17856-6, 41995-2, 4547-6, 4548-4, 4549-2, 71875-9, 96595-4 [BGP HGBA1C LOINC CODES]
  - Site-populated taxonomy DM AUDIT HGB A1C TAX
- CPT 3044F represents A1c less than ( $<$ ) 7 and will be included in the A1c less than ( $<$ ) 8 numerator.
- CPT 3046F represents A1c greater than ( $>$ ) 9 and will be included in the A1c greater than ( $>$ ) 9 numerator.
- CPT 3051F represents A1c greater than or equal to ( $\geq$ ) 7 and less than ( $<$ ) 8 and will be included in the A1c greater than or equal to ( $\geq$ ) 7 and less than ( $<$ ) 8 and A1c less than ( $<$ ) 8 numerators.

#### 2.1.2.6 GPRA 2025 Target

**Poor Glycemic Control:** During GPRA Year 2025, achieve the target rate of 12.5% for the proportion of patients with diagnosed diabetes who have poor glycemic control (defined as A1c greater than [ $>$ ] 9).

#### 2.1.2.7 Patient Lists

- List of diabetic patients with good glycemic control (A1c less than [ $<$ ] 8).
- List of diabetic patients without good Glycemic control (A1c greater than or equal to [ $\geq$ ] 8).
- List of diabetic patients with a documented A1c.
- List of diabetic patients without a documented A1c.
- List of diabetic patients with poor glycemic control (A1c greater than [ $>$ ] 9).
- List of diabetic patients with A1c greater than or equal to ( $\geq$ ) 7 and less than ( $<$ ) 8.

### 2.1.3 Diabetes: Blood Pressure Control

#### 2.1.3.1 Owner and Contact

Diabetes Program: Carmen Hardin

### 2.1.3.2 National Reporting

NATIONAL (included in IHS Performance Report; reported to OMB and Congress)

### 2.1.3.3 Denominators

1. GPRA: User Pop Diabetic patients, defined as User Population patients diagnosed with diabetes prior to the Report Period, *and* at least two visits during the Report Period, *and* two DM-related visits ever *or* DM entry on the Problem List.
2. Active Diabetic patients, defined as Active Clinical patients diagnosed with diabetes prior to the Report Period, *and* at least two visits during the Report Period, *and* two DM-related visits ever *or* DM entry on the Problem List.

### 2.1.3.4 Numerators

1. Patients with blood pressure documented during the Report Period.
2. GPRA: Patients with blood pressure less than (<) 140/90, i.e., the mean systolic value is less than (<) 140 and the mean diastolic value is less than (<) 90.

### 2.1.3.5 Definitions

#### Diabetes

First DM POV recorded in the V POV file or Problem List Entry where the status is not Deleted with Date of Onset or Date Entered prior to the Report Period:

- ICD-9: 250.00 through 250.93 or ICD-10: E10.\* through E13.\*  
[SURVEILLANCE DIABETES]
- SNOMED data set PXRDM DIABETES (Problem List only)

#### Exclusions

When calculating all BPs (using vital measurements or CPT codes), the following visits will be excluded:

- Service Category H (Hospitalization), I (In Hospital), S (Day Surgery), or O (Observation)
- Clinic code 23 (Surgical), 30 (Emergency Room [ER]), 44 (Day Surgery), 79 (Triage), C1 (Neurosurgery), or D4 (Anesthesiology)

#### BP Documented

CRS uses the mean of the last three BPs documented during the Report Period. If three BPs are not available, it uses the mean of the last two BPs, or one BP if only one is documented. If a visit contains more than one BP, the lowest BP will be used, defined as having the lowest systolic value. The mean systolic value is calculated by adding the last three (or two) systolic values and dividing by three (or two). The mean diastolic value is calculated by adding the diastolic values from the last three (or two) BPs and dividing by three (or two). If the systolic and diastolic values do not *both* meet the criteria for controlled, then the value is considered not controlled.

If CRS is not able to calculate a mean BP, it will search for CPT 0001F, 2000F, 3074F through 3080F, G9273, G9274 [BGP BP MEASURED CPT, BGP SYSTOLIC BP CPTS, BGP DIASTOLIC BP CPTS] documented during the Report Period.

### **Controlled BP**

CRS uses a mean, as described above where BP is less than (<) 140/90. If the mean systolic and diastolic values do not *both* meet the criteria for controlled, then the value is considered not controlled.

### **BP Documented and Controlled BP**

If CRS is not able to calculate a mean BP from BP measurements, it will search for the most recent of any of the following codes documented during the Report Period:

- BP Documented: CPT 0001F, 2000F, G9273, G9274 [BGP BP MEASURED CPT]; *or*
- Systolic: CPT 3074F, 3075F, or 3077F [BGP SYSTOLIC BP CPTS] WITH Diastolic: CPT 3078F, 3079F, or 3080F [BGP DIASTOLIC BP CPTS]. The systolic and diastolic values do not have to be recorded on the same day. If there are multiple values on the same day, the CPT indicating the lowest value will be used.
- The following combinations represent BP less than (<) 140/90 and will be included in the Controlled BP numerator: CPT 3074F or 3075F *and* 3078F or 3079F, *or* G9273. All other combinations will not be included in the Controlled BP numerator.

#### **2.1.3.6 GPRA 2025 Target**

During GPRA Year 2025, achieve the target rate of 57.5% for the proportion of patients with diagnosed diabetes who have achieved blood pressure control (defined as less than [ $<$ ] 140/90).



### 2.1.3.7 Patient Lists

- List of diabetic patients with controlled BP, defined as below 140/90.
- List of diabetic patients with BP that is not controlled, defined as at or above 140/90.
- List of diabetic patients who had their BP assessed.
- List of diabetic patients who did not have their BP assessed.

## 2.1.4 Statin Therapy to Reduce Cardiovascular Disease Risk in Patients with Diabetes

### 2.1.4.1 Owner and Contact

Diabetes Program: Carmen Hardin

### 2.1.4.2 National Reporting

NATIONAL (included in IHS Performance Report; reported to OMB and Congress)

### 2.1.4.3 Denominators

1. GPRA: User Pop Diabetic patients, defined as User Population patients diagnosed with diabetes prior to the Report Period, *and* at least two visits during the Report Period, *and* two DM-related visits ever *or* DM entry on the Problem List, ages 40 through 75; and User Pop Diabetic patients with documented atherosclerotic cardiovascular disease (ASCVD) or age 20 through 75 with a low-density lipoprotein (LDL) greater than or equal to ( $\geq$ ) 190 or an active diagnosis of familial hypercholesterolemia or age 40 through 75 with a 10-year ASCVD risk score of greater than or equal to ( $\geq$ ) 20%. Broken down by age groups.
2. User Pop Diabetic patients, defined as User Population patients diagnosed with diabetes prior to the Report Period, *and* at least two visits during the Report Period, *and* two DM-related visits ever *or* DM entry on the Problem List, ages 39 and under with documented ASCVD or an LDL greater than or equal to ( $\geq$ ) 190 or an active diagnosis of familial hypercholesterolemia.
3. User Pop Diabetic patients, defined as User Population patients diagnosed with diabetes prior to the Report Period, *and* at least two visits during the Report Period, *and* two DM-related visits ever *or* DM entry on the Problem List, ages 40 through 75 with documented ASCVD or an LDL greater than or equal to ( $\geq$ ) 190 or an active diagnosis of familial hypercholesterolemia or a 10-year ASCVD risk score of greater than or equal to ( $\geq$ ) 20%.

4. User Pop Diabetic patients, defined as User Population patients diagnosed with diabetes prior to the Report Period, *and* at least two visits during the Report Period, *and* two DM-related visits ever *or* DM entry on the Problem List, age 76 and older with documented ASCVD.
5. User Pop Diabetic patients, defined as User Population patients diagnosed with diabetes prior to the Report Period, *and* at least two visits during the Report Period, *and* two DM-related visits ever *or* DM entry on the Problem List, ages 40 through 75 with LDL less than ( $<$ ) 190, no ASCVD or active diagnosis of familial hypercholesterolemia, and no 10-year ASCVD risk score of greater than or equal to ( $\geq$ ) 20%.
6. User Pop Diabetic patients, defined as User Population patients diagnosed with diabetes prior to the Report Period, *and* at least two visits during the Report Period, *and* two DM-related visits ever *or* DM entry on the Problem List, ages 40 through 75; and User Pop Diabetic patients with documented ASCVD or age 20 through 75 with an LDL greater than or equal to ( $\geq$ ) 190 or an active diagnosis of familial hypercholesterolemia, or age 40 through 75 with a 10-year ASCVD risk score of greater than or equal to ( $\geq$ ) 20%, including denominator exclusions and exceptions.

#### 2.1.4.4 Numerators

1. GPRA: Patients who are statin therapy users during the Report Period or who receive an order (prescription) to receive statin therapy at any point during the Report Period.
2. Patients with any of the listed denominator exclusions or exceptions (with Denominator 6).
  - A. Patients with statin-associated muscle symptoms or an allergy to statin medication.

#### 2.1.4.5 Definitions

##### Diabetes

First DM POV recorded in the V POV file or Problem List Entry where the status is not Deleted with Date of Onset or Date Entered prior to the Report Period:

- ICD-9: 250.00 through 250.93 or ICD-10: E10.\* through E13.\*  
[SURVEILLANCE DIABETES]
- SNOMED data set PXRMI DIABETES (Problem List only)

## Atherosclerotic Cardiovascular Disease (ASCVD)

Atherosclerotic Cardiovascular Disease (ASCVD) defined as an active diagnosis of any of the following (POV or Problem List entry where the status is not Inactive or Deleted), or an ASCVD procedure (CABG, PCI, or Carotid Intervention) ever:

- Myocardial Infarction (MI) defined as any of the following:
  - Diagnosis:
    - ICD-9: 410.\*; ICD-10: I21.\* (excluding I21.A1), I22.0, I22.1, I22.8, I22.9 [BGP ECQM MI DXS]
    - SNOMED data set PXRMI BGP ECQM MI
- Ischemic Heart Disease or Other Related Diagnoses defined as any of the following:
  - Diagnosis:
    - ICD-9: 411.\*, 414.2, 414.8, 414.9; ICD-10: I23.0-I23.6, I23.8, I24.0\*, I24.8, I24.9, I25.5, I25.6, I25.82, I25.89, I25.9 [BGP ECQM ISCHEM HEART DIS DXS]
    - SNOMED data set PXRMI BGP ECQM ISCHEM HEART DIS
- Cerebrovascular Disease, Ischemic Stroke or Transient Ischemic Attack (TIA) defined as any of the following:
  - Diagnosis:
    - ICD-9: 433.\*, 434.\*, 435.0, 435.1, 435.3, 435.8, 435.9, 436, 437.1, 438.\*, V12.54; ICD-10: G45.0 through G45.2, G45.8, G45.9, G46.0 through G46.8, I63.\*, I69.\*, Z86.73 [BGP ECQM CD STROKE TIA DXS]
    - SNOMED data set PXRMI BGP ECQM CD STROKE TIA
- Atherosclerosis and Peripheral Arterial Disease defined as any of the following:
  - Diagnosis:
    - ICD-9: 414.0\*, 414.3, 414.4, 437.0, 440.\*; ICD-10: E08.51, E08.52, E09.51, E09.52, I25.1\*, I25.2, I25.5, I25.700 through I25.812, I25.82 through I25.9, I65.\*, I67.2, I70.\* [BGP ECQM ARTERIAL DISEASE DXS]
    - SNOMED data set PXRMI BGP ECQM ARTERIAL DIS
- Stable and Unstable Angina defined as any of the following:
  - Diagnosis:

- ICD-9: 411.0, 411.1, 413.0, 413.9, 429.79; ICD-10: I20.0, I20.1, I20.8, I20.9, I23.7 [BGP ECQM ANGINA DXS]
- SNOMED data set PXRMBGP ECQM ANGINA
- Coronary Artery Bypass Graft (CABG) Procedure defined as any of the following:
  - CPT 33510 through 33514, 33516 through 33519, 33521 through 33523, 33533 through 33536, S2205 through S2209
  - Procedure ICD-9: 36.1\*; ICD-10: 02100\*\*, 02103\*\*, 02104\*\*, 02110\*\*, 02113\*\*, 02114\*\*, 02120\*\*, 02123\*\*, 02124\*\*, 02130\*\*, 02133\*\*, 02134\*\* [BGP ECQM CABG PROCS]
  - SNOMED data set PXRMBGP ECQM CABG
- Percutaneous Coronary Interventions (PCI) Procedure defined as any of the following:
  - CPT 92920, 92924, 92928, 92933, 92937, 92941, 92943, 92980 (old code), 92982 (old code), 92995 (old code) [BGP PCI CPTS]
  - Procedure ICD-10: 02703\*\*, 02704\*\*, 02713\*\*, 02714\*\*, 02723\*\*, 02724\*\*, 02733\*\*, 02734\*\* [BGP ECQM PCI PROCS]
  - SNOMED data set PXRMBGP ECQM PCI
- Carotid Intervention defined as any of the following:
  - Procedure ICD-9: 00.61 through 00.65, 38.02, 38.12, 38.22, 38.30 through 38.32, 38.42, 39.22, 39.28, 88.41; ICD-10: 03\*H\*\*\*, 03\*J\*\*\*, 03\*K\*\*\*, 03\*L\*\*\*, 03\*M\*\*\*, 03\*N\*\*\*, 0G\*6\*\*\*, 0G\*7\*\*\*, 0G\*8\*\*\*, B3060ZZ, B3061ZZ, B306YZZ, B3070ZZ, B3071ZZ, B307YZZ, B3080ZZ, B3081ZZ, B308YZZ, B3160ZZ, B3161ZZ, B316YZZ, B3170ZZ, B3171ZZ, B317YZZ, B3180ZZ, B3181ZZ, B318YZZ [BGP ECQM CAROTID INTER PROCS]
  - SNOMED data set PXRMBGP ECQM CAROTID INTER

## LDL

For LDL greater than or equal to ( $\geq$ ) 190, CRS will look for any test at any time with result greater than or equal to ( $\geq$ ) 190. LDL defined as any of the following:

- LOINC taxonomy: 12773-8, 13457-7, 18261-8, 18262-6, 2089-1, 2090-9, 22748-8, 35198-1, 39469-2, 49132-4, 55440-2, 69419-0, 96258-9, 96259-7, 96597-0 [BGP LDL LOINC CODES]
- Site-populated taxonomy DM AUDIT LDL CHOLESTEROL TAX

**Familial hypercholesterolemia**

Diagnosis (POV or Problem List entry where the status is not Inactive or Deleted):

- POV ICD-10: E78.01 [BGP ECQM HYPERCHOL DXS]
- SNOMED data set PXRMBGP ECQM HYPERCHOL

**10-year ASCVD risk score**

Any of the following during the report period:

- Measurement ACC (10 YEAR ASCVD RISK).

**Denominator Exclusions**

Patients meeting any of the following conditions will be excluded from the denominator.

- Patients who are breastfeeding.
- Patients who have a diagnosis of rhabdomyolysis during the report period (POV or Problem List entry where the status is not Inactive or Deleted).

**Denominator Exceptions**

The patient is excluded from the denominator if they meet any of the following criteria and do not meet the numerator.

- Patients with statin-associated muscle symptoms or an allergy to statin medication during the Report Period.
- Patients with active liver disease or hepatic disease (including Hepatitis A or B [POV or Problem List entry where the status is not Inactive or Deleted]) or insufficiency.
- Patients who are receiving hospice or palliative care during the Report Period.
- Patients with end-stage renal disease (ESRD).
- Patients with documentation of a medical reason for not being prescribed statin therapy during the Report Period.

**Statin-associated Muscle Symptoms or an Allergy**

Defined as any of the following during the Report Period:

- POV ICD-9: 359.9, 729.1; ICD-10: G72.0, G72.9, M60.9, M79.10
- SNOMED data set PXRMBGP ECQM STATIN ADV
- POV ICD-9: 995.0 through 995.3 [BGP ASA ALLERGY 995.0-995.3] and E942.9 [BGP ADV EFF CARDIOVASC NEC]
- “Statin” or “Statins” entry (except “Nystatin”) in ART (Patient Allergies File)

- “Statin” or “Statins” (except “Nystatin”) contained within Problem List (where status is not Deleted) or in Provider Narrative field for any POV ICD-9: 995.0 through 995.3, V14.8; ICD-10: Z88.8 [BGP ASA ALLERGY 995.0-995.3 and BGP HX DRUG ALLERGY NEC]
- Problem List entry where the status is not Deleted of SNOMED data set PXRMBGPADRSTATIN
- NMI (not medically indicated) refusal for any statin at least once during the Report Period

### **Breastfeeding Definition**

Any of the following documented during the Report Period:

- The Lactation Status field in Reproductive Factors file set to “Lactating”
- Diagnosis:
  - POV ICD-10: O91.03, O91.13, O91.23, O92.03, O92.13, O92.5, O92.70, O92.79, Z39.1 [BGP ECQM BREASTFEED DXS]
  - SNOMED data set PXRMBGP ECQM BREASTFEED
- Breastfeeding Patient Education codes BF-BC, BF-BP, BF-CS, BF-EQ, BF-FU, BF-HC, BF-ON, BF-M, BF-MK, BF-N or containing SNOMED data sets PXRMBGPPTEDBREASTFEED or PXRMBGP ECQM BREASTFEED

### **Rhabdomyolysis**

- Diagnosis (POV or Problem List entry where the status is not Inactive or Deleted):
  - ICD-10: M62.82, T79.6XX\* [BGP ECQM RHABDO DXS]
  - SNOMED data set PXRMBGP ECQM RHABDO (Problem List only)

### **Active Liver Disease, Hepatic Disease or Insufficiency Definition**

Diagnosis (POV or Problem List entry where the status is not Deleted):

- POV ICD-10: B17.\*, B18.2 through B19.0, B19.20, B19.21, B19.9, K70.0 through K74.69, K75.4, O98.41\* [BGP ECQM LIVER DISEASE DXS]
- SNOMED data set PXRMBGP ECQM LIVER DIS

### **Hospice Indicator**

Hospice is defined as any of the following during the Report Period:

- SNOMED data set PXRMBGP IPC INPT ENC (Inpatient encounter) with DISCHARGE SNOMED CT data set PXRMBGP IPC DISCHG HOSPICE (discharge to home or health care facility for hospice care).

- Visit, V CPT, V POV or Problem List entry where the status is not Inactive or Deleted: SNOMED data set PXRMBGP IPC HOSPICE (hospice care ambulatory, hospice encounter, hospice diagnosis).
- CPT 99377, 99378, G0182, G9473, G9474, G9475, G9476, G9477, G9478, G9479, Q5003, Q5004, Q5005, Q5006, Q5007, Q5008, Q5010, S9126, T2042, T2043, T2044, T2045, T2046 [BGP ECQM HOSPICE CPTS].

### **Palliative Care**

Palliative care is defined as any of the following during the Report Period:

- CPT G9054
- SNOMED data set PXRMBGP ECQM PALLIATIVE ENC
- POV ICD-10: Z51.5

### **ESRD**

End Stage Renal Disease diagnosis or treatment defined as any of the following ever:

- CPT 36145 (old code), 36147, 36800, 36810, 36815, 36818, 36819, 36820, 36821, 36831 through 36833, 50300, 50320, 50340, 50360, 50365, 50370, 50380, 90951 through 90970 or old codes 90918 through 90925, 90935, 90937, 90939 (old code), 90940, 90945, 90947, 90989, 90993, 90997, 90999, 99512, 3066F, G0257, G9231, M1187, M1188, S2065, S9339 [BGP ESRD CPTS]
- Diagnosis (POV or Problem List entry where the status is not Inactive or Deleted):
  - ICD-9: 585.6, V42.0, V45.1 (old code), V45.11, V45.12, V56.\*; ICD-10: I12.0, I13.11, I13.2, N18.5, N18.6, N19., Z48.22, Z49.\*, Z91.15, Z94.0, Z99.2 [BGP ESRD PMS DXS]
  - SNOMED data set PXRMBGP END STAGE RENAL DISEASE (Problem List only)
- Procedure ICD-9: 38.95, 39.27, 39.42, 39.43, 39.53, 39.93 through 39.95, 54.98, 55.6\*; ICD-10 5A1D70Z, 5A1D80Z, 5A1D90Z [BGP ESRD PROCS]

### **Hepatitis A Definition**

Diagnosis (POV or Problem List entry where the status is not Deleted):

- ICD-10: B15.\* [BGP IPC HEP A DXS]
- SNOMED data set PXRMBGP IPC HEP A EVID

### Hepatitis B Definition

Diagnosis (POV or Problem List entry where the status is not Deleted):

- ICD-10: B16.\*, B18.0, B18.1, B19.1\*, Z22.51 [BGP ECQM HEP B DXS]
- SNOMED data set PXRMBGP ECQM HEP B

### Medical Reason for Not Being Prescribed Statin Therapy

- SNOMED data set PXRMBGP IPC NOT DONE MED for statin medication not ordered.

### Statins Numerator Logic

- **Statin Therapy Users**
  - CPT 4013F
- **Statin Medication Codes**
  - Defined with medication taxonomy BGP PQA STATIN MEDS.
  - **Statin medications and combination products:** atorvastatin (+/- amlodipine, ezetimibe), fluvastatin, lovastatin (+/- niacin), pitavastatin, pravastatin, rosuvastatin (+/- ezetimibe), simvastatin (+/- ezetimibe, niacin, sitagliptin).

Patients must have an active prescription for statin therapy during the Report Period. This includes patients who receive an order during the Report Period, or prior to the Report Period with enough days' supply to take them into the Report Period.

$$\text{Rx Days' Supply} \geq (\text{Report Period Begin Date} - \text{Prescription Date})$$

Active prescriptions include active outside medications, defined as V Med entry at any time with EHR OUTSIDE MED field not blank and the DATE DISCONTINUED field blank.

#### 2.1.4.6 GPRA 2025 Target

During GPRA Year 2025, achieve the target rate of 52.6% for the proportion of patients with diagnosed diabetes and cardiovascular disease who are on statin therapy.

#### 2.1.4.7 Patient Lists

- List of diabetic patients age 40–75 or any age with ASCVD or 20–75 years old with LDL greater than or equal to ( $\geq$ ) 190 or familial hypercholesterolemia or 40–75 years old with a 10-year ASCVD risk score greater than or equal to ( $\geq$ ) 20% with statin therapy.



- List of diabetic patients age 40–75 or any age with ASCVD or 20–75 years old with LDL greater than or equal to ( $\geq$ ) 190 or familial hypercholesterolemia or 40–75 years old with a 10-year ASCVD risk score greater than or equal to ( $\geq$ ) 20% without statin therapy.
- List of patients with denominator exclusions.

## 2.1.5 Diabetes: Nephropathy Assessment

### 2.1.5.1 Owner and Contact

Diabetes Program: Carmen Hardin

### 2.1.5.2 National Reporting

NATIONAL (included in IHS Performance Report; reported to OMB and Congress)

### 2.1.5.3 Denominators

1. GPRA: User Pop Diabetic patients, defined as User Population patients diagnosed with diabetes prior to the Report Period, *and* at least two visits during the Report Period, *and* two DM-related visits ever *or* DM entry on the Problem List.
2. Active Diabetic patients, defined as Active Clinical patients diagnosed with diabetes prior to the Report Period, *and* at least two visits during the Report Period, *and* two DM-related visits ever *or* DM entry on the Problem List.

### 2.1.5.4 Numerators

1. GPRA: Patients with nephropathy assessment, defined as an estimated Glomerular Filtration Rate (GFR) with result *and* a Quantitative Urine Albumin-to-Creatinine Ratio (UACR) during the Report Period *or* with evidence of diagnosis or treatment of end-stage renal disease (ESRD) at any time before the end of the Report Period.

### 2.1.5.5 Definitions

#### Diabetes

First DM POV recorded in the V POV file or Problem List Entry where the status is not Deleted with Date of Onset or Date Entered prior to the Report Period:

- ICD-9: 250.00 through 250.93 or ICD-10: E10.\* through E13.\*  
[SURVEILLANCE DIABETES]
- SNOMED data set PXRMI DIABETES (Problem List only)

**Estimated GFR**

- Site-populated taxonomy BGP GPRA ESTIMATED GFR TAX
- LOINC taxonomy: 33914-3, 48642-3, 48643-1, 50044-7, 50210-4, 50384-7, 62238-1, 69405-9, 70969-1, 77147-7, 88293-6, 88294-4, 94677-2, 98979-8, 98980-6, 102097-3 [BGP ESTIMATED GFR LOINC]

**Urine Albumin-to-Creatinine Ratio**

- Current Procedural Terminology (CPT) 82043 WITH 82570
- LOINC taxonomy: 14585-4, 14959-1, 30000-4, 32294-1, 77253-3, 89998-9, 9318-7 [BGP QUANT UACR LOINC]
- Site-populated taxonomy BGP QUANT UACR TESTS

**Note:** Check with your laboratory supervisor to confirm that the names added to the taxonomy reflect quantitative test values.

**ESRD**

- ESRD diagnosis or treatment defined as any of the following ever:
  - CPT 36145 (old code), 36147, 36800, 36810, 36815, 36818, 36819, 36820, 36821, 36831 through 36833, 50300, 50320, 50340, 50360, 50365, 50370, 50380, 90951 through 90970 or old codes 90918 through 90925, 90935, 90937, 90939 (old code), 90940, 90945, 90947, 90989, 90993, 90997, 90999, 99512, 3066F, G0257, G9231, M1187, M1188, S2065, S9339 [BGP ESRD CPTS]
  - Diagnosis (POV or Problem List entry where the status is not Deleted):
    - ICD-9: 585.6, V42.0, V45.1 (old code), V45.11, V45.12, V56.\*; ICD-10: I12.0, I13.11, I13.2, N18.5, N18.6, N19., Z48.22, Z49.\*, Z91.15, Z94.0, Z99.2 [BGP ESRD PMS DXS]
    - SNOMED data set PXRМ END STAGE RENAL DISEASE (Problem List only)
  - Procedure ICD-9: 38.95, 39.27, 39.42, 39.43, 39.53, 39.93 through 39.95, 54.98, 55.6\*; ICD-10 5A1D70Z, 5A1D80Z, 5A1D90Z [BGP ESRD PROCS]

**2.1.5.6 GPRA 2025 Target**

During GPRA Year 2025, achieve the target rate of 44.8% for the proportion of patients with diagnosed diabetes who are assessed for nephropathy.

**2.1.5.7 Patient Lists**

- List of diabetic patients with nephropathy assessment.

- List of diabetic patients without nephropathy assessment.

## 2.1.6 Diabetic Retinopathy

### 2.1.6.1 Owner and Contact

Gregory Smith, OD, FAAO

### 2.1.6.2 National Reporting

NATIONAL (included in IHS Performance Report; reported to OMB and Congress)

### 2.1.6.3 Denominators

1. GPRA: User Pop Diabetic patients, defined as User Population patients diagnosed with diabetes prior to the Report Period, *and* at least two visits during the Report Period, *and* two DM-related visits ever *or* DM entry on the Problem List, without a documented history of bilateral blindness or bilateral eye enucleation.
2. Active Diabetic patients, defined as Active Clinical patients diagnosed with diabetes prior to the Report Period, *and* at least two visits during the Report Period, *and* two DM-related visits ever *or* DM entry on the Problem List, without a documented history of bilateral blindness or bilateral eye enucleation.

### 2.1.6.4 Numerators

1. GPRA: Patients receiving a qualified retinal evaluation during the Report Period.

**Note:** This numerator does *not* include refusals.

- A. Patients with a validated teleretinal visit<sup>1</sup> during the Report Period.
- B. Patients with an Ophthalmology visit during the Report Period.
- C. Patients with an Optometry visit during the Report Period.

---

<sup>1</sup> Validation study properly powered and controlled against the ETDRS gold standard (American Telemedicine Association validation category 3).

### 2.1.6.5 Definitions

#### Diabetes

First DM POV recorded in the V POV file or Problem List Entry where the status is not Deleted with Date of Onset or Date Entered prior to the Report Period:

- ICD-9: 250.00 through 250.93 or ICD-10: E10.\* through E13.\*  
[SURVEILLANCE DIABETES]
- SNOMED data set PXRMI DIABETES (Problem List only)

#### Qualified Retinal Evaluation

- Diabetic retinal exam
- Other eye exam
- The following methods are qualifying for this measure:
  - Dilated retinal evaluation by an optometrist or ophthalmologist.
  - Seven standard fields stereoscopic photos (Early Treatment Diabetic Retinopathy Study [ETDRS]) evaluated by an optometrist or Ophthalmologist.
  - Any photographic method formally validated<sup>2</sup> to seven standard fields (ETDRS).

#### Diabetic Retinal Exam

Any of the following during the Report Period:

- Exam code 03 Diabetic Eye Exam (dilated retinal examination or formally validated<sup>3</sup> ETDRS photographic equivalent)
- CPT 2021F Dilated macular exam, 2022F, 2023F and G2102 Dilated retinal eye exam, 2024F, 2025F and G2103 Seven standard field stereoscopic photos with interpretation by an ophthalmologist or optometrist, 2026F, 2033F and G2104 Eye imaging formally validated<sup>4</sup> to match the diagnosis from seven standard field stereoscopic photos, S0620 Routine ophthalmological examination including refraction; new patient, S0621 Routine ophthalmological examination including refraction; established patient, S3000 Diabetic indicator; retinal eye exam, dilated, bilateral, M1220, M1221 Dilated retinal eye exam with interpretation by an ophthalmologist or optometrist or artificial intelligence. [BGP DM RETINAL EXAM CPTS].

---

<sup>2</sup> Ibid.

<sup>3</sup> Validation study properly powered and controlled against the ETDRS gold standard (American Telemedicine Association validation category 3).

<sup>4</sup> Ibid.

**Other Eye Exam**

- Non-DNKA (did not keep appointment) visits to ophthalmology or optometry clinics with an optometrist or ophthalmologist, or visits to formally validated<sup>5</sup> teleophthalmology retinal evaluation clinics. Searches for any of the following codes in the following order:
  - CPT 67028, 67038 (old code), 67039, 67040, 92002, 92004, 92012, 92014, 92018, 92019 [BGP DM EYE EXAM CPTS]
  - Clinic code A2 (Diabetic Retinopathy)<sup>6</sup>
  - Clinic codes 17<sup>7</sup> or 18<sup>8</sup> with Provider code 08, 24, or 79 or where the Service Category is not C (Chart Review) or T (Telecommunications).

**JVN Visit**

- Clinic code A2

**Ophthalmology Visit**

- Clinic code 17 with Provider code 79 where the Service Category is not C (Chart Review) or T (Telecommunications)

**Optometry Visit**

- Clinic code 18 with Provider codes 08 or 24 where the Service Category is not C (Chart Review) or T (Telecommunications)

**Bilateral Blindness**

- Diagnosis (POV or Problem List entry where the status is not Deleted):
  - ICD-9: 369.01, 369.03, 369.04; ICD-10: H54.0\* [BGP BILATERAL BLINDNESS DXS]
  - SNOMED data set PXRMBGPBILATBLINDNESS (Problem List only)
  - SNOMED data set PXRMBGPBLINDNESSUNSPECIFIED with Laterality equal to Bilateral (Problem List only)

---

<sup>5</sup> Validation study properly powered and controlled against the ETDRS gold standard (American Telemedicine Association validation category 3).

<sup>6</sup> Validated photographic (teleretinal) retinal surveillance (American Telemedicine Association validation category 3).

<sup>7</sup> Ophthalmology or Optometry clinic codes (17, 18) cannot be used for non-qualifying photographic DR examination methods<sup>5</sup> unless a dilated retinal examination by an ophthalmologist or optometrist is also accomplished during the same encounter.

<sup>8</sup> Ibid.

- One code from (SNOMED data set PXRMBGP LEFT EYE BLIND [Problem List only] *or* SNOMED data set PXRMBGP BLINDNESS UNSPECIFIED with Laterality equal to Left [Problem List only]) *and* one code from (SNOMED data set PXRMBGP RIGHT EYE BLIND [Problem List only] *or* SNOMED data set PXRMBGP BLINDNESS UNSPECIFIED with Laterality equal to Right [Problem List only])

### **Bilateral Eye Enucleation**

- CPT 65091, 65093, 65101, 65103, 65105, 65110, 65112, 65114 with modifier 50 or 09950 (50 and 09950 modifiers indicate bilateral)
- Two separate unilateral eye enucleations with visit dates at least 14 days apart with CPT 65091, 65093, 65101, 65103, 65105, 65110, 65112, 65114
- Left eye enucleation: Procedure ICD-10: 08B1\*\*\* *and* right eye enucleation: Procedure ICD-10: 08B0\*\*\* on either the same or different dates of service

### **2.1.6.6 GPRA 2025 Target**

During GPRA Year 2025, achieve the target rate of 47.6% for the proportion of patients with diagnosed diabetes who receive an annual retinal examination.

### **2.1.6.7 Patient Lists**

- List of diabetic patients who received any retinal screening.
- List of diabetic patients who did not receive any retinal screening.

## **2.2 Dental Group**

### **2.2.1 Access to Dental Services**

#### **2.2.1.1 Owner and Contact**

Dental Program: Nathan P. Mork, DDS, MPH; Timothy L. Lozon, DDS; Joel C. Knutson, DDS

#### **2.2.1.2 National Reporting**

NATIONAL (included in IHS Performance Report; reported to OMB and Congress)

#### **2.2.1.3 Denominators**

1. GPRA: User Population patients broken down by age groups (in years): 0 through 2, 3 through 5, 6 through 9, 10 through 12, 13 through 15, 16 through 21, 22 through 34, 35 through 44, 45 through 54, 55 through 74, and 75 and older.

#### 2.2.1.4 Numerators

1. GPRA: Patients with documented dental visit during the Report Period.

**Note:** This numerator does *not* include refusals.

#### 2.2.1.5 Definitions

##### Documented Dental Visit

- IHS Dental Tracking code 0000 or 0190
- RPMS Dental/ADA CDT codes 0110 through 0390, 0415 through 0471, 0601 through 0603, 0999 through 9974, 9995, 9996, 9999, D0120 through D0389, D0415 through D0470, D0701 through D0804, D0999 through D9974, D9995, D9996, D9999
- CPT code D0110 through D0390, D0415 through D9952, D9970 through D9974, D9995, D9996, D9999 [BGP DENTAL VISIT CPT CODES]
- Exam code 30
- POV ICD-10: Z01.20, Z01.21, Z13.84, Z29.3 [BGP DENTAL VISIT DXS]

#### 2.2.1.6 GPRA 2025 Target

During GPRA Year 2025, achieve the target rate of 27.0% for the proportion of patients who receive dental services.

#### 2.2.1.7 Patient Lists

- List of patients with documented dental visit.
- List of patients without documented dental visit.

### 2.2.2 Dental Sealants

#### 2.2.2.1 Owner and Contact

Dental Program: Nathan P. Mork, DDS, MPH; Timothy L. Lozon, DDS; Joel C. Knutson, DDS

#### 2.2.2.2 National Reporting

NATIONAL (included in IHS Performance Report; reported to OMB and Congress)

### 2.2.2.3 Denominators

1. GPRA: User Population patients ages 2 through 15 years. Broken down by age groups 2, 3 through 5, 6 through 9, 10 through 12, and 13 through 15.

### 2.2.2.4 Numerators

1. GPRA: Patients with at least one or more intact dental sealants.
2. Count only (no percentage comparison to denominator). For patients meeting the User Population definition, the total number of dental sealants during the Report Period. Broken down by age group 2 through 15 years.

**Note:** This numerator does *not* include refusals.

### 2.2.2.5 Definitions

#### Intact Dental Sealant

- Any of the following documented during the Report Period:
  - RPMS Dental/ADA codes 1351, 1352, 1353, D1351, D1352, D1353
  - CPT D1351, D1352, D1353
- *Or* any of the following documented during the past three years from the end of the Report Period:
  - IHS Dental Tracking code 0007

If both RPMS Dental/ADA CDT and CPT codes are found on the same visit, only the RPMS Dental/ADA CDT code will be counted. IHS Dental Tracking code 0007 will be counted regardless of whether another sealant code is submitted on the same visit or date of service.

For the count measure, only two sealants per tooth and only one repair (RPMS Dental/ADA CDT code 1353, D1353 or CPT D1353) per tooth will be counted during the Report Period. Each tooth is identified by the data element Operative Site in RPMS.

### 2.2.2.6 GPRA 2025 Target

During GPRA Year 2025, achieve the target rate of 11.8% for the proportion of patients with at least one or more intact dental sealants.

### 2.2.2.7 Patient Lists

- List of User Pop patients ages 2 through 15 years with intact dental sealant.
- List of User Pop patients ages 2 through 15 without intact dental sealant.



- Patients who received dental sealants during Report Period.

## 2.2.3 Topical Fluoride

### 2.2.3.1 Owner and Contact

Dental Program: MPH; Nathan P. Mork, DDS, MPH; Timothy L. Lozon, DDS; Joel C. Knutson, DDS

### 2.2.3.2 National Reporting

NATIONAL (included in IHS Performance Report; reported to OMB and Congress)

### 2.2.3.3 Denominators

1. GPRA: User Population patients ages 1 through 15. Broken down by age groups 1 through 2, 3 through 5, 6 through 9, 10 through 12, and 13 through 15.

### 2.2.3.4 Numerators

1. GPRA: Patients who received one or more topical fluoride applications during the Report Period.
2. Count only (no percentage comparison to denominator). For patients meeting the User Population definition, the total number of patients with at least one topical fluoride treatment during the Report Period. Broken down by age group 1 through 15.

|   |
|---|
| <b>Note:</b> This numerator does <i>not</i> include refusals. |
|---|

### 2.2.3.5 Definitions

#### Topical Fluoride Application

Defined as any of the following:

- RPMS Dental/ADA CDT codes 1201 (old code), 1203 (old code), 1204 (old code), 1205 (old code), 1206, 1208, 5986, D1206, D1208, D1354, D5986
- CPT D1201 (old code), D1203 (old code), D1204 (old code), D1205 (old code), D1206, D1208, D5986, 99188, 0792T [BGP CPT TOPICAL FLUORIDE]
- POV ICD-10: Z29.3 [BGP TOPICAL FLUORIDE DXS]

For the count measure, a maximum of one application per patient per visit is allowed.

### **2.2.3.6 GPRA 2025 Target**

During GPRA Year 2025, achieve the target rate of 27.4% for the proportion of patients who receive at least one topical fluoride application.

### **2.2.3.7 Patient Lists**

- List of User Pop patients ages 1 through 15 years with topical fluoride application.
- List of User Pop patients ages 1 through 15 without topical fluoride application.
- Patients who received at least one topical fluoride application during Report Period.

## **2.3 Immunization Group**

### **2.3.1 Influenza**

#### **2.3.1.1 Owner and Contact**

National Immunization Program: Uzo Chukwuma, MPH

#### **2.3.1.2 National Reporting**

NATIONAL (included in IHS Performance Report; reported to OMB and Congress)

#### **2.3.1.3 Denominators**

1. Active Clinical patients ages 6 months to 17 years.
2. Active Clinical patients ages 18 years and older.
3. GPRA: User Population patients ages 6 months to 17 years.
4. GPRA: User Population patients ages 18 years and older.
5. User Population patients ages 65 years and older.

#### **2.3.1.4 Numerators**

1. GPRA: Patients with influenza vaccine documented during the Report Period or with a contraindication documented at any time before the end of the Report Period.

**Note:** The only refusals included in this numerator are documented NMI refusals.

A. Patients with a contraindication or a documented NMI refusal.

### 2.3.1.5 Definitions

#### Influenza Vaccine

Any of the following documented during the Report Period:

- Immunization (CVX) codes 15, 16, 88, 111, 123, 125 through 128 135, 140, 141, 144, 149, 150, 151, 153, 155, 158, 160, 161, 166, 168, 171, 185, 186, 194, 197, 200-202, 205, 231
- POV ICD-9: V04.8 (old code), V04.81 [BGP FLU IZ DX V04.8]
- CPT 90630, 90653 through 90664, 90666, 90668, 90672 through 90674, 90682, 90685 through 90689, 90694, 90724 (old code), 90756, G0008, Q2034 through Q2039 [BGP CPT FLU]

#### Contraindication to Influenza Vaccine

Any of the following documented at any time before the end of the Report Period:

- Contraindication in the Immunization Package of “Anaphylaxis”
- PCC NMI Refusal

### 2.3.1.6 GPRA 2025 Target

Children: During GPRA Year 2025, achieve the target rate of 18.3% for the proportion of patients age 6 months through 17 years who receive an influenza immunization.

Adults: During GPRA Year 2025, achieve the target rate of 21.0% for the proportion of patients ages 18 years and older who receive an influenza immunization.

### 2.3.1.7 Patient Lists

- List of patients ages 6 months to 17 years with influenza vaccination, contraindication, or NMI refusal.
- List of patients 6 months to 17 years without influenza vaccination, contraindication, or NMI refusal.
- List of patients 18 years and older with influenza vaccination, contraindication, or NMI refusal.

- List of patients 18 years and older without influenza vaccination, contraindication, or NMI refusal.

## 2.3.2 Adult Immunizations

### 2.3.2.1 Owner and Contact

National Immunization Program: Uzo Chukwuma, MPH

### 2.3.2.2 National Reporting

NATIONAL (included in IHS Performance Report; reported to OMB and Congress)

### 2.3.2.3 Denominators

1. Active Clinical patients ages 19 years and older.
2. User Population patients ages 19 through 50.
3. User Population patients ages 51 through 65.
4. User Population patients ages 66 and older.
5. GPRA Denominator: User Population patients ages 19 and older.

### 2.3.2.4 Numerators

**Note:** The only refusals included in all numerators are documented NMI refusals.

1. Patients who have received 1 dose of Tdap/Td in the past 10 years, including contraindications (with Denominators 2 through 4).
2. Patients who have received 1 dose of Tdap ever, including contraindications (with Denominators 2 through 4).
3. Patients with influenza vaccine documented during the Report Period or with a contraindication documented at any time before the end of the Report Period (with Denominators 2 through 4).
4. Patients who have received 2 doses of Shingrix ever, including contraindications (with Denominators 3 through 4).
5. Patients with up-to-date Pneumococcal vaccine (with Denominator 4).

6. Patients who have received the 1:1:1 combination (i.e., 1 Tdap/Td in the past 10 years, 1 Tdap ever, 1 influenza during the Report Period), including contraindications (with Denominator 2).
7. Patients who have received the 1:1 combination (i.e., 1 Tdap/Td in the past 10 years, 1 Tdap ever), including contraindications (with Denominator 2).
8. Patients who have received the 1:1:1:2 combination (i.e., 1 Tdap/Td in the past 10 years, 1 Tdap ever, 1 influenza during the Report Period, 2 Shingrix), including contraindications (with Denominator 3).
9. Patients who have received the 1:1:2 combination (i.e., 1 Tdap/Td in the past 10 years, 1 Tdap ever, 2 Shingrix), including contraindications (with Denominator 3).
10. Patients who have received the 1:1:1:2:1 combination (i.e., 1 Tdap/Td in the past 10 years, 1 Tdap ever, 1 influenza during the Report Period, 2 Shingrix, 1 up-to-date Pneumococcal vaccine), including contraindications (with Denominator 4).
11. Patients who have received the 1:1:2:1 combination (i.e., 1 Tdap/Td in the past 10 years, 1 Tdap ever, 2 Shingrix, 1 up-to-date Pneumococcal vaccine), including contraindications (with Denominator 4).
12. GPRA: Patients who have received all age-appropriate immunization combinations (with Denominators 1 and 5).

### 2.3.2.5 Definitions

#### Age-appropriate Immunization Combinations

- Ages 19–50 years: 1:1 combination (i.e., 1 Tdap/Td in the past 10 years, 1 Tdap ever)
- Ages 51–65 years: 1:1:2 combination (i.e., 1 Tdap/Td in the past 10 years, 1 Tdap ever, 2 doses of Shingrix ever)
- Ages 66 years and older: 1:1:2:1 combination (i.e., 1 Tdap/Td in the past 10 years, 1 Tdap ever, 2 doses of Shingrix ever, 1 up-to-date Pneumococcal vaccine)

**Up-to-date Pneumococcal vaccine (PCV20, PCV21, PCV15, PPSV23, PCV13)** is defined as any of the following:

- Patients who have ever received PCV20 or PCV21.
- Patients who have received PCV15 followed by PPSV23 at least eight weeks apart.

- Patients who have received PPSV23 followed by PCV15, PCV20, or PCV21 at least one year apart.
- Patients who have received PCV13 at any age followed by PPSV23 at age 65 years or older at least one year apart.
- Patients with a contraindication to Pneumococcal Conjugate (PCV20, PCV21, PCV15, or PCV13) and a PPSV23 vaccine or contraindication at any time.

**Tdap Immunization:**

Any of the following documented during the applicable time frame. For pregnant patients, the Tdap must have occurred in the past 20 months and must be on or after a pregnancy visit:

- Immunization (CVX) code: 115
- CPT 90715

**Tdap Contraindication**

Any of the following documented any time before the end of the Report Period:

- Immunization Package contraindication of “Anaphylaxis”
- PCC NMI Refusal

**Td Immunization**

Any of the following documented in the past 10 years:

- Immunization (CVX) code 9, 113, 138, 139, 196
- POV ICD-9: V06.5 [BGP TD IZ DXS]
- CPT 90714, 90718

**Td Contraindication**

Any of the following documented any time before the end of the Report Period:

- Immunization Package contraindication of “Anaphylaxis”
- PCC NMI Refusal

**Influenza Vaccine**

Any of the following during the Report Period:

- Immunization (CVX) codes 15, 16, 88, 111, 123, 125 through 128, 135, 140, 141, 144, 149, 150, 151, 153, 155, 158, 160, 161, 166, 168, 171, 185, 186, 194, 197, 200-202, 205, 231
- POV ICD-9: V04.8 (old code), V04.81 [BGP FLU IZ DX V04.8]

- CPT 90630, 90653 through 90664, 90666, 90668, 90672 through 90674, 90682, 90685 through 90689, 90694, 90724 (old code), 90756, G0008, Q2034 through Q2039 [BGP CPT FLU]

**Contraindication to Influenza Vaccine**

Any of the following documented at any time before the end of the Report Period:

- Contraindication in the Immunization Package of “Anaphylaxis”
- PCC NMI Refusal

**Shingrix Vaccine**

Any of the following documented ever (the two doses must be at least 2 months apart):

- Immunization (CVX) codes 187
- CPT 90750 [BGP ZOSTER SHINGRIX CPTS]

**Contraindication to Shingrix Vaccine**

Any of the following documented at any time before the end of the Report Period:

- Contraindication in the Immunization Package of “Immune Deficiency” or “Anaphylaxis”
- PCC NMI Refusal

**Pneumococcal Polysaccharide (PPSV23) Vaccine**

Any of the following documented any time before the end of the Report Period:

- Immunization (CVX) codes 33, 109 [BGP PPSV23 CVX CODES]
- POV ICD-9: V03.82 [BGP PNEUMO IZ DXS]
- CPT 90732, G0009, G9279 [BGP PNEUMO IZ CPT DEV]

**Contraindication to Pneumococcal Polysaccharide (PPSV23) Vaccine**

Any of the following documented:

- Contraindication in the Immunization Package of “Anaphylaxis” any time before the end of the Report Period
- PCC NMI Refusal any time before the end of the Report Period

**Pneumococcal Conjugate (PCV13)**

Any of the following documented any time before the end of the Report Period:

- Immunization (CVX) codes 100, 133, 152 [BGP PCV13 CVX CODES]
- CPT 90669, 90670 [BGP PCV13 CPT CODES]

**Pneumococcal Conjugate (PCV20)**

Any of the following documented any time before the end of the Report Period:

- Immunization (CVX) codes 216 [BGP PCV20 CVX CODES]
- CPT 90677 [BGP PCV20 CPT CODES]

**Pneumococcal Conjugate (PCV21)**

Any of the following documented any time before the end of the Report Period:

- Immunization (CVX) codes 327 [BGP PCV21 CVX CODES]
- CPT 90684 [BGP PCV21 CPT CODES]

**Pneumococcal Conjugate (PCV15)**

Any of the following documented any time before the end of the Report Period:

- Immunization (CVX) codes 215 [BGP PCV15 CVX CODES]
- CPT 90671 [BGP PCV15 CPT CODES]

**Pneumococcal Conjugate Contraindication**

Any of the following documented:

- Contraindication in the Immunization Package of “Anaphylaxis” any time before the end of the Report Period
- PCC NMI Refusal any time before the end of the Report Period

**2.3.2.6 GPRA 2025 Target**

During GPRA Year 2025, achieve the target rate of 39.0% for the proportion of adult patients age 19 years and older who receive their age-appropriate immunizations.

**2.3.2.7 Patient Lists**

- List of patients ages 19–50 years with 1:1:1 combination (i.e., 1 Tdap/Td in the past 10 years, 1 Tdap ever, 1 influenza during the Report Period).
- List of patients 19–50 without 1:1:1 combination (i.e., 1 Tdap/Td in the past 10 years, 1 Tdap ever, 1 influenza during the Report Period).
- List of patients 19–50 with 1:1 combination (i.e., 1 Tdap/Td in the past 10 years, 1 Tdap ever).
- List of patients 19–50 without 1:1 combination (i.e., 1 Tdap/Td in the past 10 years, 1 Tdap ever).
- List of patients 51–65 with 1:1:1:2 combination (i.e., 1 Tdap/Td in the past 10 years, 1 Tdap ever, 1 influenza during the Report Period, 1 Zoster).



- List of patients 51–65 without 1:1:1:2 combination (i.e., 1 Tdap/Td in the past 10 years, 1 Tdap ever, 1 influenza during the Report Period, 2 Shingrix).
- List of patients 51–65 with 1:1:2 combination (i.e., 1 Tdap/Td in the past 10 years, 1 Tdap ever, 2 Shingrix).
- List of patients 51–65 without 1:1:2 combination (i.e., 1 Tdap/Td in the past 10 years, 1 Tdap ever, 2 Shingrix).
- List of patients 66 years and older with 1:1:1:2:1 combination (i.e., 1 Tdap/Td in the past 10 years, 1 Tdap ever, 1 influenza during the Report Period, 2 Shingrix, 1 up-to-date Pneumo).
- List of patients 66 years and older without 1:1:1:2:1 combination (i.e., 1 Tdap/Td in the past 10 years, 1 Tdap ever, 1 influenza during the Report Period, 2 Shingrix, 1 up-to-date Pneumo).
- List of patients 66 years and older with 1:1:2:1 combination (i.e., 1 Tdap/Td in the past 10 years, 1 Tdap ever, 2 Shingrix, 1 up-to-date Pneumo).
- List of patients 66 years and older without 1:1:2:1 combination (i.e., 1 Tdap/Td in the past 10 years, 1 Tdap ever, 2 Shingrix, 1 up-to-date Pneumo).
- List of User Population patients 19 years and older with age-appropriate immunizations.
- List of User Population patients 19 years and older without age-appropriate immunizations.

### 2.3.3 Childhood Immunizations

#### 2.3.3.1 Owner and Contact

National Immunization Program: Uzo Chukwuma, MPH

#### 2.3.3.2 National Reporting

NATIONAL (included in IHS Performance Report; reported to OMB and Congress)

#### 2.3.3.3 Denominators

1. Active Clinical patients ages 19 through 35 months at end of Report Period.
2. User Population patients active in the Immunization Package who are 19 through 35 months at the end of the Report Period.

**Note:** Sites must be running the RPMS Immunization package for this denominator. Sites not running the package will have a value of zero for this denominator.

3. GPRA: User Population patients ages 19 through 35 months at the end of the Report Period.

#### 2.3.3.4 Numerators

**Note:** *The only refusals included in the numerators in this section are documented NMI refusals.*

1. GPRA: Patients who have received the 4:3:1:3\*:3:1:4 combination (i.e., 4 DTaP, 3 Polio, 1 MMR, 3 or 4 HiB, 3 Hepatitis B, 1 Varicella, 4 Pneumococcal), including contraindications and evidence of disease.
2. Patients who have received 4 doses of DTaP ever, including contraindications.
3. Patients who have received 3 doses of Polio ever, including contraindications and evidence of disease.
4. Patients who have received 1 dose of Measles, Mumps and Rubella (MMR) vaccine ever, including contraindications, and evidence of disease.
5. Patients who have received 3 or 4 doses of Haemophilus Influenza Type B (HiB) vaccine ever, including contraindications.
6. Patients who have received 3 doses of Hepatitis B vaccine ever, including contraindications and evidence of disease.
7. Patients who have received 1 dose of Varicella vaccine ever, including contraindications and evidence of disease.
8. Patients who have received 4 doses of Pneumococcal conjugate vaccine ever, including contraindications.

#### 2.3.3.5 Definitions

##### Patient Age

Since the patient's age is calculated at the beginning of the Report Period, the age range will be adjusted to 7 through 23 months at the beginning of the Report Period, which makes the patient between the ages of 19 through 35 months at the end of the Report Period.

### Timing of Doses

Because immunization data comes from multiple sources, any immunization codes documented on dates within 10 days of each other will be considered as the same immunization.

### Active Immunization Package Patients Denominator

Same as User Population definition *except* it includes only patients flagged as active in the Immunization Package.

**Note:** Only values for the Report Period will be reported for this denominator since currently there is not a way to determine if a patient was active in the Immunization Package during the previous year or baseline periods.

### Dosage and Types of Immunizations

- 4 Doses of Diphtheria Tetanus Acellular Pertussis (DTaP) vaccine
  - 4 DTaP or Diphtheria, Tetanus and Pertussis (DTP) vaccine or Tdap
  - 1 DTaP or DTP or Tdap and 3 DT or Td
  - 1 DTaP or DTP or Tdap and 3 each of Diphtheria and Tetanus
  - 4 DT and 4 Acellular Pertussis
  - 4 Td and 4 Acellular Pertussis, or
  - 4 each of Diphtheria, Tetanus, and Acellular Pertussis
- 3 Doses of Polio
  - 3 Oral Polio Vaccine (OPV)
  - 3 Inactivated Polio Vaccine (IPV), or
  - Combination of OPV & IPV totaling 3 doses
- 1 Dose of MMR
  - MMR
  - 1 M/R and 1 Mumps
  - 1 R/M and 1 Measles, or
  - 1 each of Measles, Mumps, and Rubella
- 3 or 4 doses of HIB, depending on the vaccine administered
- 3 doses of Hep B
- 1 dose of Varicella
- 4 doses of Pneumococcal

## NMI Refusals, Contraindication, and Evidence of Disease Information

NMI refusals, evidence of disease, and contraindications for individual immunizations will also count toward meeting the definition, as defined below.

**Note:** NMI refusals are *not* counted as refusals; rather, they are counted as contraindications.

- For immunizations that allow a different number of doses (e.g., 3 or 4 HIB): To count toward the numerator with the smaller number of doses, all the patient's vaccinations must be part of the smaller dose series. For example, for a patient to count toward the HIB numerator with only 3 doses, all 3 doses must be included in the 3-dose series codes listed in the HIB definition. A patient with a mix of 3-dose and 4-dose series codes will need 4 doses to count toward the numerator. An exception to this is for the HIB vaccine: if the first 2 doses are part of the 3-dose series, then the patient only needs 3 doses (even if the third dose is included in the 4-dose series).
- Each immunization must be refused and documented separately. For example, if a patient has an NMI refusal for Rubella only, then there must be an immunization, contraindication, or separate NMI refusal for the Measles and Mumps immunizations.
- For immunizations where the required number of doses is greater than one, only one NMI refusal is necessary to be counted in the numerator. For example, if there is a single NMI refusal for Hepatitis B, the patient will be included in the numerator.
- To be counted as evidence of disease or contraindication or NMI refusal, a patient must have evidence of disease, a contraindication, or an NMI refusal for any of the immunizations in the numerator. For example, if a patient was Rubella immune but had a Measles and Mumps immunization, the patient would be counted as having evidence of disease for MMR.
- For immunizations where the required number of doses is greater than one, only one contraindication is necessary to be counted in the numerator. For example, if there is a single contraindication for HiB, the patient will be included in the numerator.
- Evidence of disease will be checked for at any time in the child's life (prior to the end of the Report Period).

### NMI Refusal Definitions

PCC Refusal type NMI for any of the following codes:

- DTaP
  - Immunization (CVX) codes 20, 50, 102, 106, 107, 110, 120, 130, 132, 146

- CPT 90696 through 90698, 90700, 90721, 90723 [BGP CPT DTAP/DTP/TDAP]
- DTP
  - Immunization (CVX) codes 1, 22, 102, 198
  - CPT 90701, 90711 (old code), 90720 [BGP CPT DTAP/DTP/TDAP]
- Tdap (Tetanus, Diphtheria and Acellular Pertussis) vaccine
  - Immunization (CVX) code 115
  - CPT 90715 [BGP CPT DTAP/DTP/TDAP]
- DT
  - Immunization (CVX) code 28
  - CPT 90702
- Td
  - Immunization (CVX) codes 9, 113, 138, 139, 196
  - CPT 90714, 90718
- Diphtheria
  - CPT 90719
- Tetanus
  - Immunization (CVX) codes 35, 112
  - CPT 90703
- Acellular Pertussis
  - Immunization (CVX) code 11
- OPV
  - Immunization (CVX) codes 2, 89
  - CPT 90712
- IPV
  - Immunization (CVX) codes 10, 89, 110, 120, 130, 132, 146
  - CPT 90696 through 90698, 90711 (old code), 90713, 90723
- MMR
  - Immunization (CVX) codes 3, 94
  - CPT 90707, 90710
- M/R (Measles and Rubella)
  - Immunization (CVX) code 4
  - CPT 90708

- R/M (Rubella/Mumps)
  - Immunization (CVX) code 38
  - CPT 90709 (old code)
- Measles
  - Immunization (CVX) code 5
  - CPT 90705
- Mumps
  - Immunization (CVX) code 7
  - CPT 90704
- Rubella
  - Immunization (CVX) code 6
  - CPT 90706
- HiB
  - Immunization (CVX) codes 17, 22, 46 through 49, 50, 51, 102, 120, 132, 146, 148, 198
  - CPT 90644 through 90648, 90697, 90698, 90720, 90721, 90737 (old code), 90748 [BGP HIB CPT]
- Hepatitis B
  - Immunization (CVX) codes 8, 42 through 45, 51, 102, 104, 110, 132, 146, 189, 193, 198, 220
  - CPT 90636, 90697, 90723, 90731 (old code), 90739, 90740, 90743 through 90748, 90759, G0010 [BGP HEPATITIS CPTS]
- Varicella
  - Immunization (CVX) codes 21, 94
  - CPT 90710, 90716
- Pneumococcal
  - Immunization (CVX) codes 33, 100, 109, 133, 152, 215, 216, 327 [BGP CHILD IZ PNEUMO CVX CODES]
  - CPT 90669, 90670, 90671, 90677, 90684, 90732, G0009, G9279 [BGP CHILD IZ PNEUMO CPT CODES]

### **Contraindication Definitions**

- Encephalopathy due to vaccination
  - POV or Problem List entry where the status is not Deleted:
    - ICD-9: 323.51; ICD-10: G04.32 [BGP ENCEPHALOPATHY DXS]

- SNOMED data set PXRMM BGP IPC IZ ENCEPHAL
- Vaccine adverse-effect  
POV or Problem List entry where the status is not Deleted: ICD-9: E948.4 through E948.6; ICD-10: T50.A15\* [BGP VACCINE ADVERSE EFFECT]
- Immunodeficiency  
POV or Problem List entry where the status is not Deleted:
  - ICD-9: 279.\*; ICD-10: D80.\*, D81.0 through D81.7, D81.89, D81.9, D82.\* through D84.\*, D89.3, D89.8\*, D89.9 [BGP IMMUNODEFICIENCY DXS]
  - SNOMED data set PXRMM BGP IPC IMMUNE DIS
- HIV  
POV or Problem List entry where the status is not Deleted:
  - ICD-9: 042, 079.53, V08; ICD-10: B20, B97.35, Z21, O98.711 through O98.73 [BGP HIV/AIDS DXS]
  - SNOMED data set PXRMM BGP IPC HIV (Problem List only)
- Lymphoreticular cancer, multiple myeloma, or leukemia  
POV or Problem List entry where the status is not Deleted:
  - ICD-9: 200.00 through 208.92; ICD-10: C81.00 through C86.6, C88.2 through C88.9, C90.00 through C93.\*, C94.00 through C94.32, C94.80, C95.\*, C96.0 through C96.4, C96.9, C96.A, C96.Z [BGP LYMPHO CANCER DXS]
  - SNOMED data set PXRMM BGP IPC LYMPH CANCER

### Immunization Definitions

- DTaP
  - Immunization (CVX) codes 20, 50, 102, 106, 107, 110, 120, 130, 132, 146
  - POV ICD-9: V06.1
  - CPT 90696 through 90698, 90700, 90721, 90723 [BGP CPT DTAP/DTP/TDAP]
- DTaP Contraindication Definition
  - Immunization Package contraindication of “Anaphylaxis”
  - Encephalopathy due to vaccination with a vaccine adverse-effect
- DTP
  - Immunization (CVX) codes 1, 22, 102, 198
  - POV ICD-9: V06.1, V06.2, V06.3 [BGP DTP IZ DXS]

- CPT 90701, 90711 (old code), 90720 [BGP CPT DTAP/DTP/TDAP]
- DTP Contraindication Definition
  - Immunization Package contraindication of “Anaphylaxis”
- Tdap
  - Immunization (CVX) code 115
  - CPT 90715 [BGP CPT DTAP/DTP/TDAP]
- Tdap Contraindication Definition
  - Immunization Package contraindication of “Anaphylaxis”
- DT
  - Immunization (CVX) code 28
  - POV ICD-9: V06.5 [BGP TD IZ DXS]
  - CPT 90702
- DT Contraindication Definition
  - Immunization Package contraindication of “Anaphylaxis”
- Td
  - Immunization (CVX) codes 9, 113, 138, 139, 196
  - POV ICD-9: V06.5 [BGP TD IZ DXS]
  - CPT 90714, 90718 [BGP CPT TDAP/TD]
- Td Contraindication Definition
  - Immunization Package contraindication of “Anaphylaxis”
- Diphtheria
  - POV ICD-9: V03.5 [BGP DIPHTHERIA IZ DXS]
  - CPT 90719
- Diphtheria Contraindication Definition
  - Immunization Package contraindication of “Anaphylaxis”
- Tetanus
  - Immunization (CVX) codes 35, 112
  - POV ICD-9: V03.7 [BGP TETANUS TOXOID IZ DXS]
  - CPT 90703
- Tetanus Contraindication Definition
  - Immunization Package contraindication of “Anaphylaxis”
- Acellular Pertussis
  - Immunization (CVX) code 11



- POV ICD-9: V03.6 [BGP PERTUSSIS IZ DXS]
- Acellular Pertussis Contraindication Definition
  - Immunization Package contraindication of “Anaphylaxis”
- OPV
  - Immunization (CVX) codes 2, 89
  - CPT 90712
- OPV Contraindication Definition
  - Immunization Package contraindication of “Immune Deficiency”
- IPV
  - Immunization (CVX) codes 10, 89, 110, 120, 130, 132, 146
  - POV ICD-9: V04.0, V06.3 [BGP IPV IZ DXS]
  - CPT 90696 through 90698, 90711 (old code), 90713, 90723
- IPV Evidence of Disease Definitions
  - POV or PCC Problem List (active or inactive) ICD-9: 730.70 through 730.79; ICD-10: M89.6\* [BGP OPV EVID DISEASE]
  - SNOMED data set PXRMBGP POLIO (Problem List only)
- IPV Contraindication Definition
  - Immunization Package contraindication of “Anaphylaxis” or “Neomycin Allergy”
- MMR
  - Immunization (CVX) codes 3, 94
  - POV ICD-9: V06.4 [BGP MMR IZ DXS]
  - CPT 90707, 90710
- MMR Contraindication Definitions
  - Immunization Package contraindication of “Anaphylaxis,” “Immune Deficiency,” or “Neomycin Allergy”
  - Immunodeficiency
  - HIV
  - Lymphoreticular cancer, multiple myeloma or leukemia
- M/R
  - Immunization (CVX) code 4
  - CPT 90708
- M/R Contraindication Definition
  - Immunization Package contraindication of “Anaphylaxis”

- R/M
  - Immunization (CVX) code 38
  - CPT 90709 (old code)
- R/M Contraindication Definition
  - Immunization Package contraindication of “Anaphylaxis”
- Measles
  - Immunization (CVX) code 5
  - POV ICD-9: V04.2 [BGP MEASLES IZ DXS]
  - CPT 90705
- Measles Evidence of Disease Definition
  - POV or PCC Problem List (active or inactive) ICD-9: 055\*; ICD-10: B05.\* [BGP MEASLES EVIDENCE]
  - SNOMED data set PXRMBGP IPC MEASLES EVID (Problem List only)
- Measles Contraindication Definition
  - Immunization Package contraindication of “Anaphylaxis”
- Mumps
  - Immunization (CVX) code 7
  - POV ICD-9: V04.6 [BGP MUMPS IZ DXS]
  - CPT 90704
- Mumps Evidence of Disease Definition
  - POV or PCC Problem List (active or inactive) ICD-9: 072\*; ICD-10: B26.\* [BGP MUMPS EVIDENCE]
  - SNOMED data set PXRMBGP IPC MUMPS EVID (Problem List only)
- Mumps Contraindication Definition
  - Immunization Package contraindication of “Anaphylaxis”
- Rubella
  - Immunization (CVX) code 6
  - POV ICD-9: V04.3 [BGP RUBELLA IZ DXS]
  - CPT 90706
- Rubella Evidence of Disease Definitions
  - POV or PCC Problem List (active or inactive) ICD-9: 056\*, 771.0; ICD-10: B06.\* [BGP RUBELLA EVIDENCE]

- SNOMED data set PXRMBGP IPC RUBELLA EVID (Problem List only)
- Rubella Contraindication Definition
  - Immunization Package contraindication of “Anaphylaxis”
- HiB
  - 3-dose series:
    - Immunization (CVX) codes 49, 51
    - CPT 90647, 90748
  - 4-dose series:
    - Immunization (CVX) codes 17, 22, 46 through 48, 50, 102, 120, 132, 146, 148, 198
    - POV ICD-9: V03.81 [BGP HIB IZ DXS]
    - CPT 90644 through 90646, 90648, 90697, 90698, 90720, 90721, 90737 (old code)
- HiB Contraindication Definition
  - Immunization Package contraindication of “Anaphylaxis”
- Hepatitis B
  - Immunization (CVX) codes 8, 42 through 45, 51, 102, 104, 110, 132, 146, 189, 193, 198, 220
  - CPT 90636, 90697, 90723, 90731 (old code), 90739, 90740, 90743 through 90748, 90759, G0010 [BGP HEPATITIS CPTS]
- Hepatitis B evidence of disease definition
  - POV or PCC Problem List (active or inactive) ICD-10: B16.\*, B18.0, B18.1, B19.1\* [BGP HEP EVIDENCE]
  - SNOMED data set PXRMBGP IPC HEP B EVID (Problem List only)
- Hepatitis B contraindication definition
  - Immunization Package contraindication of “Anaphylaxis”
- Varicella
  - Immunization (CVX) codes 21, 94
  - POV ICD-9: V05.4 [BGP VARICELLA IZ DXS]
  - CPT 90710, 90716
- Varicella Evidence of Disease Definition
  - POV or PCC Problem List (active or inactive) ICD-9: 052\*, 053\*; ICD-10: B01.\* through B02.\* [BGP VARICELLA EVIDENCE]

- SNOMED data set PXRMBGP IPC VZV EVID (Problem List only)
- Immunization Package contraindication of “Hx of Chicken Pox” or “Immune”
- Varicella Contraindication Definitions
  - Immunization Package contraindication of “Anaphylaxis,” “Immune Deficiency,” or “Neomycin Allergy”
  - Immunodeficiency
  - HIV
  - Lymphoreticular cancer, multiple myeloma, or leukemia
- Pneumococcal
  - Immunization (CVX) codes 33, 100, 109, 133, 152, 215, 216, 327 [BGP CHILD IZ PNEUMO CVX CODES]
  - POV ICD-9: V03.82 [BGP PNEUMO IZ DXS]
  - CPT 90669, 90670, 90671, 90677, 90684, 90732, G0009, G9279 [BGP CHILD IZ PNEUMO CPT CODES]
- Pneumococcal Contraindication Definition
  - Immunization Package contraindication of “Anaphylaxis”

### 2.3.3.6 GPRA 2025 Target

During GPRA Year 2025, achieve the target rate of 37.8% for the proportion of AI/AN children ages 19 through 35 months who have received the recommended immunizations.

**Notes:** In FY 2013, the GPRA measure changed to the 4:3:1:3\*:3:1:4 combination, which includes 3 or 4 HiB.

In FY 2011, the GPRA measure changed to the 4:3:1:3:3:1:4 combination, which includes pneumococcal.

### 2.3.3.7 Patient Lists

**Note:** Because age is calculated at the beginning of the Report Period, the patient’s age on the list will be between 7 and 23 months.

- List of User Population patients ages 19 through 35 months who received the 4:3:1:3\*:3:1:4 combination (4 DTaP, 3 Polio, 1 MMR, 3 or 4 HiB, 3 Hep B, 1 Varicella, 4 Pneumococcal).

- List of User Population patients ages 19 through 35 months who have not received the 4:3:1:3\*:3:1:4 combination (4 DTaP, 3 Polio, 1 MMR, 3 or 4 HiB, 3 Hep B, 1 Varicella, 4 Pneumococcal). If a patient did not have all doses in a multiple dose vaccine, the IZ will not be listed. For example, if a patient only had 2 DTaP, no IZ will be listed for DTaP.
- List of Active Immunization Package patients ages 19 through 35 months who received the 4:3:1:3\*:3:1:4 combination (4 DTaP, 3 Polio, 1 MMR, 3 or 4 HiB, 3 Hep B, 1 Varicella, and 4 Pneumococcal).
- List of Active Immunization Package patients ages 19 through 35 months who have not received the 4:3:1:3\*:3:1:4 combination (4 DTaP, 3 Polio, 1 MMR, 3 or 4 HiB, 3 Hep B, 1 Varicella, and 4 Pneumococcal). If a patient did not have all doses in a multiple dose vaccine, the IZ will not be listed. For example, if a patient only had 2 DTaP, no IZ will be listed for DTaP.

## 2.4 Cancer Screening Group

### 2.4.1 Cervical Cancer Screening

#### 2.4.1.1 Owner and Contact

CAPT Suzanne England, DNP, APRN

#### 2.4.1.2 National Reporting

NATIONAL (included in IHS Performance Report; reported to OMB and Congress)

#### 2.4.1.3 Denominators

1. Female Active Clinical patients ages 24 through 64 without a documented history of hysterectomy.

**Note:** Patients must be at least 24 years of age at the beginning of the Report Period and less than 65 years of age as of the end of the Report Period.

2. GPRA: Female User Population patients ages 24 through 64 without a documented history of hysterectomy.

**Note:** Patients must be at least 24 years of age at the beginning of the Report Period and less than 65 years of age as of the end of the Report Period.

3. Female User Population patients ages 24 through 29 without documented history of hysterectomy.
4. Female User Population patients ages 30 through 64 without documented history of hysterectomy.

#### 2.4.1.4 Numerators

**Note:** The numerators in this section do *not* include refusals.

1. GPRA: Patients with a Pap smear documented in the past three years, or if patient is 30–64 years old, either a Pap Smear documented in the past three years or a Pap Smear and an HPV DNA documented on the same day in the past five years or HPV Primary in the past five years.
2. Patients with a Pap Smear documented in the past three years (with Denominators 3 and 4).
3. Patients with a Pap Smear documented three to five years ago, and an HPV DNA documented on the same day in the past five years (with Denominator 4).
4. Patients with HPV Primary in the past 5 years (with Denominator 4).

#### 2.4.1.5 Definitions

##### Age

Age of the patient is calculated at the beginning of the Report Period. Patients must be at least 24 years of age at the beginning of the Report Period and less than 65 years of age as of the end of the Report Period.

##### Hysterectomy

Defined as any of the following ever:

- Procedure ICD-9: 68.4 through 68.8; ICD-10: 0UTC\*ZZ, 0UT90ZL, 0UT9\*ZZ [BGP HYSTERECTOMY PROCEDURES]
- CPT 51925, 56308 (old code), 58150, 57540, 57545, 57550, 57555, 57556, 58152, 58200 through 58294, 58548, 58550 through 58554, 58570 through 58573, 58575, 58951, 58953 through 58954, 58956, 59135 [BGP HYSTERECTOMY CPTS]
- Diagnosis (POV or Problem List entry where the status is not Deleted):
  - ICD-9: 618.5, 752.43, V88.01, V88.03; ICD-10: N99.3, Z12.72, Z90.710, Z90.712, Q51.5 [BGP HYSTERECTOMY DXS]

- SNOMED data set PXR M BGP HYSTERECTOMY DX (Problem List only)

- Women’s Health procedure called Hysterectomy

### Pap Smear

- Lab PAP SMEAR
- POV ICD-9: V76.2 Screen Mal Neop-Cervix, V72.32 Encounter for Pap Cervical Smear to Confirm Findings of Recent Normal Smear Following Initial Abnormal Smear, 795.0\*; ICD-10: R87.61\*, R87.810, R87.820, Z01.42, Z12.4 [BGP PAP SMEAR DXS]
- CPT 88141 through 88154, 88160 through 88167, 88174 through 88175, G0123, G0124, G0141, G0143 through G0145, G0147, G0148, P3000, P3001, Q0091 Screening Pap Smear [BGP CPT PAP]
- Women’s Health procedure called Pap Smear and where the result does not have “ERROR/DISREGARD”
- LOINC taxonomy: 10524-7, 18500-9, 19762-4, 19763-2, 19764-0, 19765-7, 19766-5, 19767-3, 19768-1, 19769-9, 19770-7, 19771-5, 19772-3, 19773-1, 19774-9, 33717-0, 39086-4, 47527-7, 47528-5, 49034-2, 49050-8, 104866-9 [BGP PAP LOINC CODES]
- Site-populated taxonomy BGP GPRA PAP SMEAR TAX

### HPV DNA

**Note:** CRS will only search for a documented HPV DNA if the patient had a Pap Smear three to five years ago.

- Lab HPV
- POV ICD-9: V73.81, 079.4, 795.05, 795.09, 795.15, 795.19, 796.75, 796.79; ICD-10: B97.7, R85.618, R85.81, R85.82, R87.628, R87.810, R87.811, R87.820, R87.821, Z11.51 [BGP HPV DXS]
- CPT 87620 through 87622 (old codes), 87623 through 87625, G0476, 0429U [BGP HPV CPTS]
- Women’s Health procedure called HPV Screen and where the result does *not* have “ERROR/DISREGARD”
- Women’s Health procedure called Pap Smear and where the HPV field equals Yes
- LOINC taxonomy: 10705-2, 11083-3, 11481-9, 12222-6, 12223-4, 13321-5, 13322-3, 14499-8, 14500-3, 14501-1, 14502-9, 14503-7, 14504-5, 14505-2, 14506-0, 16280-0, 17398-9, 17399-7, 17400-3, 17401-1, 17402-9, 17403-7, 17404-5, 17405-2, 17406-0, 17407-8, 17408-6, 17409-4, 17410-2, 17411-0,

17412-8, 18478-8, 18479-6, 18480-4, 21440-3, 21441-1, 22434-5, 30167-1, 32047-3, 38372-9, 42481-2, 42770-8, 43170-0, 43209-6, 43210-4, 43211-2, 44543-7, 44544-5, 44545-2, 44546-0, 44547-8, 44548-6, 44549-4, 44550-2, 44551-0, 48560-7, 49891-5, 49896-4, 50642-8, 55298-4, 55299-2, 56140-7, 59263-4, 59264-2, 59420-0, 61372-9, 61373-7, 61374-5, 61375-2, 61376-0, 61377-8, 61378-6, 61379-4, 61380-2, 61381-0, 61382-8, 61383-6, 61384-4, 61385-1, 61386-9, 61387-7, 61388-5, 61389-3, 61390-1, 61391-9, 61392-7, 61393-5, 61394-3, 61395-0, 61396-8, 6510-2, 6511-0, 6512-8, 6513-6, 6514-4, 6515-1, 6516-9, 69002-4, 69358-0, 70061-7, 71431-1, 71432-9, 73732-0, 73959-9, 74763-4, 74776-6, 74777-4, 74778-2, 74779-0, 75406-9, 75664-3, 75694-0, 75695-7, 75696-5, 76498-5, 77375-4, 77376-2, 77377-0, 77378-8, 77379-6, 77380-4, 77394-5, 77395-2, 77396-0, 77399-4, 77400-0, 7975-6, 81439-2, 82354-2, 82456-5, 82457-3, 82675-0, 86560-0, 86561-8, 86562-6, 86563-4, 86564-2, 91073-7, 91851-6, 91852-4, 91853-2, 91854-0, 91855-7, 91856-5, 93777-1, 93778-9, 94425-6, 95539-3, 95538-5, 95537-7, 95536-9, 95535-1, 95534-4, 95533-6, 95532-8, 97156-4, 97157-2, 101822-5, 104132-6, 104170-6, 104752-1, 104766-1, 106051-6 [BGP HPV LOINC CODES]

- Site-populated taxonomy BGP HPV TAX

#### **HPV Primary**

- Lab HPV Primary
- CPT 87624
- LOINC taxonomy: 59420-0, 71432-9, 77377-0, 77379-6, 77378-8, 95532-8, 97156-4 [BGP HPV PRIMARY LOINC CODES]
- Site-populated taxonomy BGP HPV PRIMARY TAX

#### **2.4.1.6 GPRA 2025 Target**

During GPRA Year 2025, achieve the target rate of 35.6% for the proportion of female patients ages 24 through 64 years without a documented history of hysterectomy who have had a Pap screen within the previous three years, or if the patient is over 30, have had a Pap screen in the past three years or a Pap screen and HPV DNA on the same day within the previous five years.

#### **2.4.1.7 Patient Lists**

- List of female patients with a Pap smear documented in the past three years or Pap plus HPV in past five years.
- List of female patients without a Pap smear documented in the past three years or Pap plus HPV in past five years.



## 2.4.2 Cancer Screening: Mammogram Rates

### 2.4.2.1 Owner and Contact

CAPT Suzanne England, DNP, APRN

### 2.4.2.2 National Reporting

NATIONAL (included in IHS Performance Report; reported to OMB and Congress)

### 2.4.2.3 Denominators

**Note:** For both definitions in this section, the patients must be at least 52 years of age as of the beginning of the Report Period and less than 75 years of age as of the end of the Report Period.

1. Female Active Clinical patients ages 52 through 74 years, without a documented bilateral mastectomy or two separate unilateral mastectomies.
2. GPRA: Female User Population patients ages 52 through 74 years, without a documented bilateral mastectomy or two separate unilateral mastectomies.

### 2.4.2.4 Numerators

1. GPRA: All patients who had a Mammogram documented in the past two years.

**Note:** This numerator does *not* include refusals.

### 2.4.2.5 Definitions

#### Age

Age of the patient is calculated at the beginning of the Report Period. Patients must be at least 52 years old at of the beginning of the Report Period and younger than 75 years of age as of the end of the Report Period.

#### Bilateral Mastectomy

- CPT M1280, 19300.50 through 19307.50 *or* 19300 through 19307 with modifier 09950 (50 and 09950 modifiers indicate bilateral), or old codes 19180, 19200, 19220, or 19240, with modifier of 50 or 09950 [BGP MASTECTOMY CPTS]
- Procedure ICD-9: 85.42, 85.44, 85.46, 85.48; ICD-10: 0HBV0ZZ, 0HCV0ZZ, 0HDV0ZZ, 0HTV0ZZ [BGP MASTECTOMY PROCEDURES]

- Diagnosis (POV or Problem List entry where the status is not Deleted):
  - ICD-10: Z90.13 [BGP MASTECTOMY DXS]
  - SNOMED data set PXRMBGP BILAT MASTECTOMY (Problem List only)

### **Two Separate Unilateral Mastectomies**

Requires either of the following:

- Must have one code that indicates a right mastectomy and one code that indicates a left mastectomy
- Must have two separate occurrences on two different dates of service for one code that indicates a mastectomy on unknown side and one code that indicates either a right or left mastectomy, or two codes that indicate a mastectomy on unknown side

### **Right Mastectomy**

- Diagnosis (POV or Problem List entry where the status is not Deleted):
  - ICD-10: Z90.11 [BGP RIGHT MASTECTOMY DXS]
  - SNOMED data set PXRMBGP RIGHT MASTECTOMY (Problem List only)
- Procedure ICD-10: 07T50ZZ, 07T80ZZ, 0HBT0ZZ, 0HCT0ZZ, 0HDT0ZZ, 0HTT0ZZ [BGP UNI RIGHT MASTECTOMY PROCS]

### **Left Mastectomy**

- Diagnosis (POV or Problem List entry where the status is not Deleted):
  - ICD-10: Z90.12 [BGP LEFT MASTECTOMY DXS]
  - SNOMED data set PXRMBGP LEFT MASTECTOMY (Problem List only)
- Procedure ICD-10: 07T60ZZ, 07T90ZZ, 0HBU0ZZ, 0HCU0ZZ, 0HDU0ZZ, 0HTU0ZZ [BGP UNI LEFT MASTECTOMY PROCS]

### **Mastectomy on Unknown Side**

- CPT 19300 through 19307, or old codes 19180, 19200, 19220, 19240 [BGP UNI MASTECTOMY CPTS]
- Procedure ICD-9: 85.41, 85.43, 85.45, 85.47 [BGP UNI MASTECTOMY PROCEDURES]

### **Mammogram**

- Radiology or CPT 77046 through 77049, 77052 through 77059, 77061 through 77063, 77065 through 77067, 76090 (old code), 76091 (old code), 76092 (old code), G0206, G0204, G0202, G0279 [BGP CPT MAMMOGRAM]

- POV ICD-9: V76.11, V76.12, 793.80 Abnormal mammogram, unspecified, 793.81 Mammographic microcalcification, 793.89, Other abnormal findings on radiological exam of breast; ICD-10: R92.0, R92.1, R92.8, Z12.31 [BGP MAMMOGRAM DXS]
- Procedure ICD-9: 87.36, 87.37; ICD-10: BH00ZZZ, BH01ZZZ, BH02ZZZ [BGP MAMMOGRAM PROCEDURES]
- Women’s Health procedure called Mammogram Screening, Mammogram Dx Bilat, Mammogram Dx Unilat and where the mammogram result does *not* have “ERROR/DISREGARD”

#### 2.4.2.6 GPRA 2025 Target

During GPRA Year 2025, achieve the target rate of 40.5% for the proportion of female patients ages 52 through 74 years who have had mammography screening within the last two years.

#### 2.4.2.7 Patient Lists

- List of female patients with a Mammogram documented in the past two years.
- List of female patients without a Mammogram documented in the past two years.

### 2.4.3 Colorectal Cancer Screening

**Note:** Numerator does not include Double Contrast Barium Enema (DCBE).

#### 2.4.3.1 Owner and Contact

Epidemiology Program: Don Haverkamp

#### 2.4.3.2 National Reporting

NATIONAL (included in IHS Performance Report; reported to OMB and Congress)

#### 2.4.3.3 Denominators

**Note:** For this section, the patient’s age at the beginning of the Report Period must be at least 45 years of age and 46 years of age at the end of the Report Period.

1. Active Clinical patients ages 45 through 75 without a documented history of colorectal cancer or total colectomy.

2. GPRA: User Population patients ages 45 through 75 without a documented history of colorectal cancer or total colectomy.

#### 2.4.3.4 Numerators

1. GPRA: Patients who have had any Colorectal Cancer (CRC) screening, defined as any of the following:
  - A. Fecal Occult Blood Test (FOBT) or FIT during the Report Period
  - B. Flexible sigmoidoscopy or CT colonography in the past five years
  - C. Colonoscopy in the past 10 years
  - D. FIT-DNA in the past three years
2. Patients with Fecal Occult Blood test (FOBT) or Fecal Immunochemical Test (FIT) during the Report Period.

#### 2.4.3.5 Definitions

##### Denominator Exclusions

Any diagnosis ever of one of the following:

- Colorectal Cancer
  - Diagnosis (POV or Problem List entry where the status is not Deleted):
    - ICD-9: 153.\*, 154.0, 154.1, 197.5, V10.05, V10.06; ICD-10: C18.\*, C19, C20, C21.2, C21.8, C78.5, Z85.030, Z85.038, Z85.048 [BGP COLORECTAL CANCER DXS]
    - SNOMED data set PXR COLORECTAL CANCER (Problem List only)
- Total Colectomy
  - CPT 44150 through 44151, 44152 (old code), 44153 (old code), 44155 through 44158, 44210 through 44212, M1295 [BGP TOTAL COLECTOMY CPTS]
  - Procedure ICD-9: 45.8\*; ICD-10: 0DTE\*ZZ [BGP TOTAL COLECTOMY PROCS]

##### Colorectal Cancer Screening

The most recent of any of the following during applicable time frames:

- FOBT or FIT
  - CPT 82270, 82274, 89205 (old code), G0328 [BGP FOBT CPTS]

- LOINC taxonomy: 12503-9, 12504-7, 14563-1, 14564-9, 14565-6, 2335-8, 27396-1, 27401-9, 27925-7, 27926-5, 29771-3, 42912-6, 42913-4, 56490-6, 56491-4, 57905-2, 58453-2, 50196-5, 57803-9, 80372-6, 104738-0 [BGP FOBT LOINC CODES]
- Site-populated taxonomy BGP GPRA FOB TESTS
- Flexible Sigmoidoscopy
  - Procedure ICD-9: 45.24; ICD-10: 0DJD8ZZ [BGP SIG PROCS]
  - CPT 45330 through 45347, 45349, 45350, G0104 [BGP SIG CPTS]
- CT Colonography
  - CPT 74261 through 74263 [BGP CT COLONOGRAPHY CPTS]
- Colonoscopy
  - Procedure ICD-9: 45.22, 45.23, 45.25, 45.42, 45.43; ICD-10: 0D5E4ZZ, 0D5E8ZZ, 0D5F4ZZ, 0D5F8ZZ, 0D5G4ZZ, 0D5G8ZZ, 0D5H4ZZ, 0D5H8ZZ, 0D5K4ZZ, 0D5K8ZZ, 0D5L4ZZ, 0D5L8ZZ, 0D5M4ZZ, 0D5M8ZZ, 0D5N4ZZ, 0D5N8ZZ, 0D9E3ZX, 0D9E4ZX, 0D9E7ZX, 0D9E8ZX, 0D9F3ZX, 0D9F4ZX, 0D9F7ZX, 0D9F8ZX, 0D9G3ZX, 0D9G4ZX, 0D9G7ZX, 0D9G8ZX, 0D9H3ZX, 0D9H4ZX, 0D9H7ZX, 0D9H8ZX, 0D9K3ZX, 0D9K4ZX, 0D9K7ZX, 0D9K8ZX, 0D9L3ZX, 0D9L4ZX, 0D9L7ZX, 0D9L8ZX, 0D9M3ZX, 0D9M4ZX, 0D9M7ZX, 0D9M8ZX, 0D9N3ZX, 0D9N4ZX, 0D9N7ZX, 0D9N8ZX, 0DBE3ZX, 0DBE4ZX, 0DBE7ZX, 0DBE8ZX, 0DBE8ZZ, 0DBF3ZX, 0DBF4ZX, 0DBF7ZX, 0DBF8ZX, 0DBF8ZZ, 0DBG3ZX, 0DBG4ZX, 0DBG7ZX, 0DBG8ZX, 0DBG8ZZ, 0DBH3ZX, 0DBH4ZX, 0DBH7ZX, 0DBH8ZX, 0DBH8ZZ, 0DBK3ZX, 0DBK4ZX, 0DBK7ZX, 0DBK8ZX, 0DBK8ZZ, 0DBL3ZX, 0DBL4ZX, 0DBL7ZX, 0DBL8ZX, 0DBL8ZZ, 0DBM3ZX, 0DBM4ZX, 0DBM7ZX, 0DBM8ZX, 0DBM8ZZ, 0DBN3ZX, 0DBN4ZX, 0DBN7ZX, 0DBN8ZX, 0DBN8ZZ, 0DJD8ZZ [BGP COLO PROCS]
  - CPT 44388 through 44394, 44397, 44401 through 44408, 45355, 45378 through 45393, 45398, G0105, G0121, G2204, G9252, G9253 [BGP COLO CPTS]
- FIT-DNA
  - CPT 81528, G0464 [BGP FIT-DNA CPTS]
  - LOINC taxonomy: 77353-1, 77354-9 [BGP FIT-DNA LOINC CODES]
  - Site-populated taxonomy BGP FIT-DNA TESTS

#### 2.4.3.6 GPRA 2025 Target

During GPRA Year 2025, achieve the target rate of 24.6% for the proportion of clinically appropriate patients ages 45 through 75 years who have received colorectal screening.

### 2.4.3.7 Patient Lists

- List of patients ages 45 through 75 years with CRC screening.
- List of patients ages 45 through 75 years without CRC screening.

## 2.4.4 Tobacco Use and Exposure Assessment

### 2.4.4.1 Owner and Contact

Chris Lamer, PharmD

Epidemiology Program: Dayle Knutson, RN, BSN

### 2.4.4.2 National Reporting

NATIONAL (included in IHS Performance Report; *not* reported to OMB and Congress)

### 2.4.4.3 Denominators

1. User Population patients ages 5 years and older.

### 2.4.4.4 Numerators

1. Patients screened for tobacco use during the Report Period.
2. Patients identified during the Report Period as current tobacco users.
  - A. Current smokers
  - B. Current smokeless tobacco users
  - C. Current E-Cigarette users
3. Patients identified as smokers of substances other than tobacco during the Report Period.
4. Patients exposed to environmental tobacco smoke (ETS) during the Report Period.

### 2.4.4.5 Definitions

#### Tobacco Screening

At least one of the following:

- Any Health Factor for category Tobacco [C004], TOBACCO (SMOKING) [C017], TOBACCO (SMOKELESS – CHEWING/DIP) [C016], E-CIGARETTES [C019], TOBACCO (EXPOSURE) [C015]

- POV or Problem List entry where the status is not Inactive or Deleted:
  - ICD-10: F17.2\*, O99.33\*, Z71.6, Z72.0, Z87.891 [BGP TOBACCO DXS]
  - SNOMED data set PXRMBGP TOBACCO SCREENED (Problem List only)
- RPMS Dental/ADA CDT code 1320, D1320
- Patient Education codes containing “TO-”, “-TO”, “-SHS,” 305.1, 305.1\* (old codes), 649.00 through 649.04, V15.82, F17.2\*, O99.33\*, Z71.6, Z72.0, Z87.891, D1320, 99406, 99407, G0030, 1034F, 1035F, 1036F, 1000F, 4000F, 4001F, 4004F, G9016, G9275, G9276, G9458, or SNOMED 408939007 or data set PXRMBGP TOBACCO SCREENED
- CPT D1320, 99406, 99407, G0030, G9016, G9275, G9276, G9458, 1034F (Current Tobacco Smoker), 1035F (Current Smokeless Tobacco User), 1036F (Current Tobacco Non-User), 1000F (Tobacco Use Assessed), 4000F, 4001F, 4004F [BGP TOBACCO SCREEN CPTS]

### **Tobacco Users**

- Health Factors: Current Smokeless [F003]; Current Smoker and Smokeless [F030]; Current E-cigarette user w/nicotine [F124]; Current Smoker, status unknown [F002]; Current smoker, every day [F108]; Current smoker, some day [F109]; Heavy Tobacco Smoker [F121]; Light Tobacco Smoker [F122]
- Diagnosis (POV or Problem List entry where the status is not Inactive or Deleted):
  - ICD-10: F17.2\*0, F17.2\*3, F17.2\*8, F17.2\*9, O99.33\*, Z71.6, Z72.0 [BGP TOBACCO USER DXS]
  - SNOMED data set PXRMBGP CURRENT TOBACCO (Problem List only)
- CPT 99406, 99407, 1034F, 1035F, 4000F, 4001F, G9016, G9276, G9458 [BGP TOBACCO USER CPTS]

### **Current Smokers**

- Health Factors: Current Smoker and Smokeless [F030]; Current Smoker, status unknown [F002]; Current smoker, every day [F108]; Current smoker, some day [F109]; Heavy Tobacco Smoker [F121]; Light Tobacco Smoker [F122]
- Diagnosis (POV or Problem List entry where the status is not Inactive or Deleted):
  - ICD-10: F17.200, F17.203 through F17.210, F17.213 through F17.219, F17.290, F17.293 through F17.299, O99.33\* [BGP GPRA SMOKING DXS]

- SNOMED data set PXRMBGP TOBACCO SMOKER (Problem List only)
- CPT 99406, 99407, G9016, 1034F [BGP SMOKER CPTS]

### **Current Smokeless**

- Health Factors: Current Smokeless [F003], Current Smoker and Smokeless [F030]
- Diagnosis (POV or Problem List entry where the status is not Inactive or Deleted):
  - ICD-10: F17.220, F17.223 through F17.229 [BGP GPRA SMOKELESS DXS]
  - SNOMED data set PXRMBGP TOBACCO SMOKELESS (Problem List only)
- CPT 1035F [BGP SMOKELESS TOBACCO CPTS]

### **E-Cigarettes**

- Health Factors: Current E-cigarette user w/nicotine [F124]

### **Other substance**

- Health Factors: Current E-cig user w/other substance(s) [F155]

### **ETS**

Health Factors: Smoker in Home [F006], Exposure to Environmental Tobacco Smoke [F031]

#### **2.4.4.6 Patient Lists**

- List of patients with documented tobacco screening.
- List of patients without documented tobacco screening.
- List of patients identified as current tobacco users, both smokers and smokeless users.

#### **2.4.5 Tobacco Cessation**

##### **2.4.5.1 Owner and Contact**

Chris Lamer, PharmD

Epidemiology Program: Dayle Knutson, RN BSN

##### **2.4.5.2 National Reporting**

NATIONAL (included in IHS Performance Report; reported to OMB and Congress)



### 2.4.5.3 Denominators

1. GPRA: User Population patients identified as current tobacco users. Broken down by sex and age groups: younger than 12 years old, 12 through 17, 18 and older.
2. Active clinical patients identified as current tobacco users.

### 2.4.5.4 Numerators

1. Patients who have received tobacco cessation counseling or received a prescription for a smoking cessation aid anytime during the Report Period.
2. Patients identified as having quit their tobacco use anytime during the Report Period.
3. GPRA: Patients who received tobacco cessation counseling, received a prescription for a tobacco cessation aid, or quit their tobacco use anytime during the Report Period.

### 2.4.5.5 Definitions

#### Denominator

Current Tobacco Users or Tobacco Users in Cessation:

CRS will search first for all health factors documented in the TOBACCO (SMOKING) [C017], TOBACCO (SMOKELESS – CHEWING/DIP) [C016] and E-CIGARETTES [C019] categories during the Report Period.

If health factor(s) are found and at least one of them is one of the health factors listed below, the patient is counted as a current tobacco user. The patient is not counted as receiving cessation counseling.

Tobacco User Health Factors (TUHF):

- SMOKER category:
  - Current Smoker, status unknown [F002]
  - Current Smoker, every day [F108]
  - Current Smoker, some day [F109]
  - Heavy Tobacco Smoker [F121]
  - Light Tobacco Smoker [F122]
- SMOKELESS category:
  - Current Smokeless [F003]
- E-Cigarettes category:
  - Current E-cigarette user w/nicotine [F124]

If no TUHF listed above was found in each of the three categories (SMOKER, SMOKELESS, and E-Cigarettes) during the specified time frame, CRS will then search for the most recent health factor in each category documented during an *expanded* time frame of any time prior to the Report Period. For example, a patient with the most recent health factor being documented five years prior to the Report Period.

**Note:** If multiple health factors of the same category were documented on the same date and if any of them are TUHF(s), all the health factors of the same category will be considered as TUHF(s).

If a health factor is found during the expanded time frame and is a TUHF, the patient will be considered a current tobacco user.

If a TUHF is not found, the patient is considered a non-tobacco user and will not be included in the denominator.

A patient is considered a smoker if any of the following health factors are found: Current Smoker, status unknown [F002]; Current Smoker, every day [F108]; Current Smoker, some day [F109]; Heavy Tobacco Smoker [F121]; Light Tobacco Smoker [F122].

A patient is considered a smokeless user if any of the following health factors are found: Current Smokeless [F003].

A patient is considered an E-Cigarette user if any of the following health factors are found: Current E-cigarette user w/nicotine [F124].

### **Tobacco Cessation Counseling**

Any of the following documented anytime during the Report Period:

- Patient education codes containing “TO-,” “-TO,” “-SHS,” 305.1, 305.1\* (old codes), 649.00 through 649.04, V15.82, F17.2\*, O99.33\*, Z71.6, Z72.0, Z87.891, D1320, 99406, 99407, 4000F, G0030, G9016, G9458, or SNOMED data set PXRMBGP TOBACCO SCREENED.
- Clinic code 94 (tobacco cessation clinic)
- RPMS Dental/ADA CDT code 1320, D1320
- CPT D1320, 99406, 99407, G0030, G9016, G9458, 4000F
- POV ICD-10: Z71.6

### Prescription for Tobacco Cessation Aid

Any of the following documented anytime during the Report Period:

- Prescription for medication in the site-populated BGP ECQM TOB CESSATION MEDS taxonomy that does not have a comment of RETURNED TO STOCK
- Prescription for any medication with name containing “NICOTINE PATCH,” “NICOTINE POLACRILEX,” “NICOTINE INHALER,” “NICOTINE NASAL SPRAY” that does not have a comment of RETURNED TO STOCK
- CPT 4001F

### Quit Tobacco Use

In order to meet the Quit Tobacco Use numerator, the patient must quit all forms of tobacco for which s/he is a user.

If the patient is a smoker, the last documented health factor must be one of the following: Previous (former) smoker [F004], Cessation-Smoker [F024].

If the patient is a smokeless user, the last documented health factor must be one of the following: Previous (former) smokeless [F005], Cessation-Smokeless [F023].

If the patient is an E-Cigarette user, the last documented health factor must be one of the following: Former E-cigarette user [F126], Cessation ENDS user [F125].

### Health Factor to SNOMED Mapping

IHS Tobacco Health Factor codes are mapped to SNOMED CT codes. These codes that are represented by the Health Factors are listed in Table 2-1, Table 2-2, and Table 2-3.

Table 2-1: Smoking Health Factors mapped to SNOMED codes

| Health Factor                  | SNOMED CT       |
|--------------------------------|-----------------|
| Current Smoker, status unknown | 77176002        |
| Current Smoker, every day      | 449868002       |
| Current Smoker, some day       | 428041000124106 |
| Heavy Tobacco Smoker           | N/A             |
| Light Tobacco Smoker           | N/A             |
| Previous (former) Smoker       | 8517006         |
| Cessation-Smoker               | 8517006         |

Table 2-2: Smokeless Tobacco Health Factors mapped to SNOMED codes

| Health Factor               | SNOMED CT |
|-----------------------------|-----------|
| Current Smokeless           | N/A       |
| Previous (former) Smokeless | N/A       |
| Cessation-Smokeless         | N/A       |

Table 2-3: E-Cigarette Health Factors mapped to SNOMED codes

| Health Factor                       | SNOMED CT |
|-------------------------------------|-----------|
| Current E-Cigarette User w/nicotine | 785889008 |
| Former E-Cigarette User             | N/A       |
| Cessation ENDS User                 | N/A       |

#### 2.4.5.6 GPRA 2025 Target

During GPRA Year 2025, achieve the target rate of 27.5% for the proportion of tobacco-using patients who receive tobacco cessation intervention.

#### 2.4.5.7 Patient Lists

- List of tobacco users with documented tobacco cessation intervention or who quit their tobacco use.
- List of tobacco users without documented tobacco cessation intervention and did not quit their tobacco use.
- List of tobacco users with documented tobacco cessation intervention.
- List of tobacco users without documented tobacco cessation intervention.
- List of tobacco users who quit tobacco use.
- List of tobacco users who did not quit tobacco use.

## 2.5 Behavioral Health Group

### 2.5.1 Alcohol Screening

#### 2.5.1.1 Owner and Contact

Kateri Fletcher-Sahmaunt, IHS Division of Behavioral Health (DBH)

### 2.5.1.2 National Reporting

NATIONAL (included in IHS Performance Report; reported to OMB and Congress)

### 2.5.1.3 Denominators

1. Active Clinical Plus BH patients ages 9 through 75 years.
2. Female Active Clinical patients ages 14 through 46 (child-bearing age).
3. Female Active Clinical patients ages 14 to 46 screened for alcohol use during the Report Period, not including refusals.

**Note:** This denominator does *not* include patients with a screening refusal or an alcohol-related diagnosis or patient education.

4. GPRA: User Population patients ages 9 through 75.
5. User Population patients ages 9 through 75 screened for alcohol use during the Report Period, not including refusals or patient education.

**Note:** This denominator does *not* include patients with a screening refusal or an alcohol-related diagnosis or patient education.

### 2.5.1.4 Numerators

1. GPRA: Patients screened for alcohol use or who had an alcohol-related diagnosis during the Report Period (with Denominators 1, 4).

**Note:** This numerator does *not* include refusals or alcohol-related patient education.

2. Patients with alcohol-related patient education during the Report Period (with Denominator 4).
3. Patients who screened positive for alcohol use (with Denominators 3 and 5).
4. Patients screened for alcohol use, had an alcohol-related diagnosis, or received alcohol-related patient education during the Report Period (with Denominator 2).

**Note:** This numerator does *not* include refusals.

### 2.5.1.5 Definitions

**Note:** Visit data can be found in PCC or BHS.

#### Alcohol Screening

Any of the following during the Report Period:

- PCC Exam code 35
- Health Factors:
  - Any Cut Down, Annoyed, Guilty, and Eye Opener (CAGE) Health Factor [F018-F022] or CAGE-Adapted to Include Drugs (CAGE-AID) [F210-F214]
  - Any TAPS-Alcohol [F175-F179]
- Behavioral Health System (BHS) problem code 29.1 (screening for alcoholism)
- CPT 99408, 99409, G0396, G0397, G0442, G0443, G2011, G2196, G2197, H0049, H0050, 3016F [BGP ALCOHOL SCREENING CPTS]
- Measurement of AUDT, AUDC, or CRFT

#### Alcohol-Related Diagnosis

Any of the following during the Report Period:

- Alcohol-related Diagnosis
  - POV or Current Problem List ICD-10: F10.1\*, F10.20, F10.220 through F10.29, F10.920 through F10.982, F10.99, G62.1 [BGP ALCOHOL DXS]
  - SNOMED data set PXRMBGP ETOH RELATED DX (Problem List only)
  - BHS POV or Problem Codes 10, 12.1, 14.2, 17.1, 18.1, 20.1, 22.1, 27, 29

#### Alcohol-Related Patient Education

Any of the following during the Report Period:

- All Patient Education codes containing “AOD-” or “-AOD,” “CD-” or “-CD” (old codes), or F10.1\*, F10.20, F10.220-F10.29, F10.920-F10.982, F10.99, G62.1 [BGP ALCOHOL DXS], 99408, 99409, G0396, G0397, G0442, G0443, G2011, G2196, G2197, H0049, H0050, 3016F [BGP ALCOHOL SCREENING CPTS], or SNOMED 408947007 or data set PXRMBGP ETOH RELATED DX.

#### Positive Screen for Alcohol Use

Any of the following for patients with alcohol screening:

- Exam code 35 Alcohol Screening result of “Positive”

- Health factor of CAGE or CAGE-AID result of 1/4, 2/4, 3/4, or 4/4 [F019-F022], [F211-F214]
- TAPS-Alcohol health factors Alcohol-Problem Use [F176], Alcohol-High Risk [F177], or Alcohol-Early Remission, but at Risk [F179]
- CPT G0396, G0397, G2011, G2197, 99408, 99409 [BGP ALCOHOL POSITIVE SCRIN CPTS]
- AUDT result of greater than or equal ( $\geq$ ) to 8, AUDC result of greater than or equal to ( $\geq$ ) 4 for men and greater than or equal to ( $\geq$ ) 3 for women, CRFT result of 2–6

### 2.5.1.6 GPRA 2025 Target

During GPRA Year 2025, achieve the target rate of 36.0% for the proportion of patients ages 9 through 75 years who receive screening for alcohol use.

### 2.5.1.7 Patient Lists

- List of female Active Clinical patients ages 14–46 years with documented screening.
- List of female Active Clinical patients ages 14–46 years without documented screening.
- List of female Active Clinical patients 14–46 with a positive alcohol screen.
- List of female Active Clinical patients 14–46 with a negative alcohol screen/no result.
- List of User Population patients 9–75 with documented alcohol screening.
- List of User Population patients 9–75 without documented alcohol screening.
- List of patients 9–75 with a positive alcohol screen.
- List of patients 9–75 with a negative alcohol screen/no result.

## 2.5.2 Screening, Brief Intervention, and Referral to Treatment (SBIRT)

### 2.5.2.1 Owner and Contact

Kateri Fletcher-Sahmaunt, IHS Division of Behavioral Health (DBH)

### 2.5.2.2 National Reporting

NATIONAL (included in National GPRA/GPRAMA Report; reported to OMB and Congress)

### 2.5.2.3 Denominators

1. Active Clinical Plus BH patients age 9 through 75 years who screened positive for risky or harmful alcohol use during the Report Period.
2. User Population patients age 9 through 75 years.
3. GPRA: User Population patients age 9 through 75 years who screened positive for risky or harmful alcohol use during the Report Period.

### 2.5.2.4 Numerators

1. Patients screened at an Ambulatory or Telemedicine visit for risky or harmful alcohol use (with Denominator 1).
  - A. Patients screened positive for risky or harmful alcohol use.
  - B. Patients provided a brief negotiated interview (BNI) or Brief Intervention (BI) at an Ambulatory or Telemedicine visit within seven days of screen.
2. GPRA: Patients provided a Brief Negotiated Interview (BNI) or Brief Intervention (BI) at an Ambulatory or Telemedicine visit within seven days of screen (with Denominators 2 and 3).
  - A. Patients who received a BNI/BI on same day as screen.
  - B. Patients who received a BNI/BI one to three days after screen.
  - C. Patients who received a BNI/BI four to seven days after screen.
  - D. Patients who were referred treatment within seven days of screen.

### 2.5.2.5 Definitions

**Note:** Visit data can be found in PCC or BHS.

#### Ambulatory Care

- Service Category A (Ambulatory)

#### Telemedicine Visit

- Service Category M (Telemedicine)

#### Screening for Risky or Harmful Alcohol Use

Any of the following:

- Exam code 35
- Health Factors:
  - Any CAGE [F018-F022] or CAGE-AID [F210-F214]



- Any TAPS-Alcohol [F175-F179]
- CPT G0396, G0397, G0442, G0443, G2011, G2196, G2197, H0049, H0050, 99408, 99409, 3016F [BGP ALCOHOL SCREENING CPTS]
- Measurement in PCC of AUDT, AUDC, CRFT

### **Positive Screen for Risky or Harmful Alcohol Use**

CRS will look for the most recent positive screen during the Report Period, if any. A positive screen is defined as any of the following for the screening performed:

- Exam code 35 Alcohol Screening result of Positive
- Health factor of CAGE or CAGE-AID result of 1/4, 2/4, 3/4, or 4/4 [F019-F022, F211-F214]
- TAPS-Alcohol Health Factors: Alcohol-Problem Use [F176], Alcohol-High Risk [F177], or Alcohol-Early Remission, but at Risk [F179]
- Any of the following:
  - AUDT result greater than or equal to ( $\geq$ ) 8
  - AUDC result greater than or equal to ( $\geq$ ) 4 (men)
  - AUDC result greater than or equal to ( $\geq$ ) 3 (women)
  - CRFT result greater than or equal to ( $\geq$ ) 2 and CRFT result less than or equal to ( $\leq$ ) 6

### **BNI/BI**

Any of the following documented at the Ambulatory Care or Telemedicine visit or within seven days of the Ambulatory Care or Telemedicine visit, which excludes chart reviews:

- CPT G0396, G0397, G2011, G2200, H0050, 99408, 99409, 96150 through 96155 [BGP BNI CPTS]
- Patient education code containing AOD-BNI, G0396, G0397, G2011, G2200, H0050, 99408, 99409, 96150 through 96155, or SNOMED code 408947007

### **Referral to Treatment**

- Patient education code AOD-TX

#### **2.5.2.6 GPRA 2025 Target**

During GPRA Year 2025, achieve the target rate of 15.8% for the proportion of patients ages 9 through 75 years who screened positive for risky or harmful alcohol use and who received a Brief Negotiated Interview (BNI) or Brief Intervention (BI) within seven days of screen.

### 2.5.2.7 Patient Lists

- List of UP patients ages 9–75 years who were screened for risky or harmful alcohol use.
- List of UP patients ages 9–75 years who were not screened for risky or harmful alcohol use.
- List of UP patients ages 9–75 who screened positive for risky or harmful alcohol use.
- List of UP patients ages 9–75 who received a BNI/BI within seven days of screen.
- List of UP patients ages 9–75 who did not receive a BNI/BI within seven days of screen.
- List of UP patients ages 9–75 who screened positive for risky or harmful alcohol use and who were referred treatment within seven days of screen.
- List of UP patients ages 9–75 who screened positive for risky or harmful alcohol use and who were not referred treatment within seven days of screen.
- List of UP patients ages 9–75 who screened positive for risky or harmful alcohol use and who received a BNI/BI within seven days of screen.
- List of UP patients ages 9–75 who screened positive for risky or harmful alcohol use and who did not receive a BNI/BI within seven days of screen.

## 2.5.3 Intimate Partner (Domestic) Violence Screening

### 2.5.3.1 Owner and Contact

IHS Division of Behavioral Health (DBH): Tamara D. James, PhD; Nicole Stahlmann, MN, RN, SANE-A, AFN-BC, FNE-A/P

### 2.5.3.2 National Reporting

NATIONAL (included in IHS Performance Report; reported to OMB and Congress)

### 2.5.3.3 Denominators

1. Female Active Clinical patients ages 14 through 46 years.
2. GPRAMA: Female User Population patients ages 14 through 46 years.

### 2.5.3.4 Numerators

1. GPRAMA: Patients screened for intimate partner (domestic) violence (IPV/DV) at any time during the Report Period.

**Note:** This numerator does *not* include refusals.

- A. Patients with documented IPV/DV exam.
- B. Patients with IPV/DV-related diagnosis.
- C. Patients provided with education or counseling about IPV/DV.

### 2.5.3.5 Definitions

**Note:** Visit data can be found in PCC or BHS.

#### Intimate Partner Violence/Domestic Violence (IPV/DV) Screening

Defined as one of the following:

- IPV/DV Exam
  - Exam code 34
  - BHS IPV/DV exam
- IPV/DV Related Diagnosis
  - POV or Problem List ICD-10: T74.11X\*, T74.21X\*, T74.31X\*, T74.91X\*, T76.11X\*, T76.21X\*, T76.31X\*, T76.91X\*, Z91.410 [BGP DV DXS]
  - SNOMED data set PXRMBGP IPV DV DX (Problem List only)
  - BHS POV 43.\*, 44.\*
- IPV/DV Patient Education
  - Patient Education codes containing “DV-” or “-DV,” T74.11XA, T74.21XA, T74.31XA, T74.91XA, T76.11XA, T76.21XA, T76.31XA, T76.91XA, Z91.410 [BGP IPV/DV EDUC DXS], or SNOMED 413457006 or data set PXRMBGP IPV DV DX
- IPV/DV Counseling
  - POV ICD-10: Z69.11 [BGP IPV/DV COUNSELING ICDS]

### 2.5.3.6 GPRA 2025 Target

During GPRA Year 2025, achieve the target rate of 30.5% for the proportion of female patients ages 14 through 46 years who receive screening for domestic violence.

### 2.5.3.7 Patient Lists

- List of female patients 14 through 46 years with documented IPV/DV screening.

- List of female patients 14 through 46 years without documented IPV/DV screening.

## 2.5.4 Depression Screening

### 2.5.4.1 Owner and Contact

Pamela End of Horn, IHS Division of Behavioral Health (DBH)

### 2.5.4.2 National Reporting

NATIONAL (included in IHS Performance Report; reported to OMB and Congress)

### 2.5.4.3 Denominators

1. Active Clinical patients 12 through 17 years old.
2. Active Clinical patients ages 18 years and older.
3. GPRA: User Population patients ages 12 through 17 years. Broken down by sex.
4. GPRA: User Population patients ages 18 years and older. Broken down by sex.
5. Active Diabetic patients, defined as Active Clinical patients diagnosed with diabetes prior to the Report Period, *and* at least two visits during the Report Period, *and* two DM-related visits ever *or* DM entry on the Problem List. Broken down by sex.

### 2.5.4.4 Numerators

1. GPRA: Patients screened for depression or diagnosed with a mood disorder at any time during the Report Period.

**Note:** This numerator does *not* include refusals.

- A. Patients screened for depression during the Report Period.
- B. Patients with a diagnosis of a mood disorder during the Report Period.
- C. Patients who were screened in a Behavioral Health clinic.

### 2.5.4.5 Definitions

**Note:** Visit data can be found in PCC or BHS.

## Diabetes

First DM POV recorded in the V POV file or Problem List Entry where the status is not Deleted with Date of Onset or Date Entered prior to the Report Period:

- ICD-9: 250.00 through 250.93 or ICD-10: E10.\* through E13.\* [SURVEILLANCE DIABETES]
- SNOMED data set PXRMI DIABETES (Problem List only)

## Depression Screening

Any of the following:

- Exam code 36
- POV ICD-10: Z13.3\* [BGP DEPRESSION SCRNI DXS]
- CPT 1220F, 3725F, G0444 [BGP DEPRESSION SCREEN CPTS]
- BHS problem code 14.1 (screening for depression)
- Measurement of PHQ9, PHQT, or EPDS.

## Mood Disorders

At least two visits during the Report Period with POV for: Major Depressive Disorder, Dysthymic Disorder, Depressive Disorder Not Otherwise Specified (NOS), Bipolar I or II Disorder, Cyclothymic Disorder, Bipolar Disorder NOS, Mood Disorder Due to a General Medical Condition, Substance-induced Mood Disorder, or Mood Disorder NOS. These POV codes are:

- ICD-10: F01.51, F06.31 through F06.34, F1\*.\*4, F10.159, F10.180, F10.181, F10.188, F10.259, F10.280, F10.281, F10.288, F10.959, F10.980, F10.981, F10.988, F30.\*, F31.0 through F31.71, F31.73 through F31.75, F31.77, F31.81 through F31.9, F32.\* through F39, F43.21, F43.23 [BGP MOOD DISORDERS]
- BHS POV 14, 15

## Behavioral Health Clinic

- Clinic codes C4, C9, 14, 43, 48

### 2.5.4.6 GPRA 2025 Target

Ages 12–17 years: During GPRA Year 2025, achieve the target rate of 36.1% for the proportion of patients ages 12 through 17 years who receive annual screening for depression.

Ages 18 years and older: During GPRA Year 2025, achieve the target rate of 39.6% for the proportion of adults ages 18 and older who receive annual screening for depression.

### 2.5.4.7 Patient Lists

- List of User Population patients 18 years and older screened for depression or diagnosed with mood disorder.
- List of User Population patients 18 years and older *not* screened for depression or diagnosed with mood disorder.

## 2.6 Cardiovascular Disease Related Group

### 2.6.1 Childhood Weight Control

#### 2.6.1.1 Owner and Contact

Dr. Thomas Faber

#### 2.6.1.2 National Reporting

NATIONAL (included in IHS Performance Report; reported to OMB and Congress)

#### 2.6.1.3 Denominators

1. Active Clinical patients ages 2 through 5 years for whom a BMI could be calculated.
2. GPRA: User Population patients ages 2 through 5 years for whom a BMI could be calculated.

#### 2.6.1.4 Numerators

1. GPRA: Patients with a BMI at or above the 95<sup>th</sup> percentile.

#### 2.6.1.5 Definitions

##### Age

All patients who are between the ages of 2 and 5 years at the beginning of the Report Period and who do not turn age 6 during the Report Period are included in this measure. Age in the age groups is calculated based on the date of the most current BMI found. For example, a patient may be 2 at the beginning of the time period but is 3 at the time of the most current BMI found. That patient will fall into the Age 3 group.

## BMI

CRS looks for the most recent BMI in the Report Period. CRS calculates the BMI at the time the report is run, using the National Health and Nutrition Examination Survey (NHANES) II. Height and weight measurements must be taken on the same day at any time during the Report Period. The BMI values for this measure are reported differently than in an Obesity Assessment since this age group is for children ages 2 through 6 years, whose BMI values are age dependent. The BMI values are categorized as Overweight for patients with a BMI in the 85th through 94th percentile and Obese for patients with a BMI at or above the 95th percentile.

Patients whose BMI is either greater or lesser than the Data Check Limit range, as shown in Table 2-4, will not be included in the report counts for Overweight or Obese.

Table 2-4: NHANES II BMI

| Low-High Ages | Sex    | BMI (Overweight) | BMI (Obese) | Data Check Limits BMI > | Data Check Limits BMI < |
|---------------|--------|------------------|-------------|-------------------------|-------------------------|
| 2-2           | Male   | 17.7             | 18.7        | 36.8                    | 7.2                     |
| 2-2           | Female | 17.5             | 18.6        | 37.0                    | 7.1                     |
| 3-3           | Male   | 17.1             | 18.0        | 35.6                    | 7.1                     |
| 3-3           | Female | 17.0             | 18.1        | 35.4                    | 6.8                     |
| 4-4           | Male   | 16.8             | 17.8        | 36.2                    | 7.0                     |
| 4-4           | Female | 16.7             | 18.1        | 36.0                    | 6.9                     |
| 5-5           | Male   | 16.9             | 18.1        | 36.0                    | 6.9                     |
| 5-5           | Female | 16.9             | 18.5        | 39.2                    | 6.8                     |

### 2.6.1.6 GPRA 2025 Target

During GPRA Year 2025, achieve the target rate of 22.0% for the proportion of children with a BMI of 95% or higher.

### 2.6.1.7 Patient Lists

List of patients ages 2 through 5 years with a BMI at or above the 95th percentile.

## 2.6.2 Controlling High Blood Pressure – Million Hearts

### 2.6.2.1 Owner and Contact

Dr. Dena Wilson; Chris Lamer, PharmD

### 2.6.2.2 National Reporting

NATIONAL (included in IHS Performance Report; reported to OMB and Congress)

### 2.6.2.3 Denominators

1. GPRA: Million Hearts (National Quality Forum [NQF] 0018): User Population patients ages 18 through 85 years diagnosed with hypertension and no documented history of ESRD or current diagnosis of pregnancy.
2. User Population patients ages 18 through 59 years diagnosed with hypertension and no documented history of ESRD or current diagnosis of pregnancy.
3. User Population patients ages 60 through 85 years diagnosed with hypertension and no documented history of ESRD or current diagnosis of pregnancy.

### 2.6.2.4 Numerators

1. GPRA: Million Hearts (NQF 0018): Patients with blood pressure less than (<) 140/90, i.e., the systolic value is less than (<) 140 *and* the diastolic value is less than (<) 90 (with Denominator 1).
2. Patients with blood pressure less than (<) 140/90 (i.e., the systolic value is less than 140 *and* the diastolic value is less than 90) (with Denominator 2).
3. Patients with blood pressure less than (<) 150/90 (i.e., the systolic value is less than 150 *and* the diastolic value is less than 90) (with Denominator 3).

### 2.6.2.5 Definitions

#### Age

Age of the patient is calculated as of the end of the Report Period.

#### Hypertension

Diagnosis (POV or Problem List entry where the status is not Inactive or Deleted):

- ICD-9: 401.\*; ICD-10: I10 [BGP HYPERTENSION DXS] during the Report Period or the year prior to the Report Period.
- SNOMED data set PXRMESSENTIAL HYPERTENSION (Problem List only).



**ESRD**

Any of the following ever:

- CPT 36145 (old code), 36147, 36800, 36810, 36815, 36818, 36819, 36820, 36821, 36831 through 36833, 50300, 50320, 50340, 50360, 50365, 50370, 50380, 90918 through 90925 (old codes), 90935, 90937, 90939 (old code), 90940, 90945, 90947, 90951 through 90970, 90989, 90993, 90997, 90999, 99512, 3066F, G0257, G9231, M1187, M1188, S2065, S9339 [BGP ESRD CPTS]
- Diagnosis (POV or Problem List entry where the status is not Deleted):
  - ICD-9: 585.6, V42.0, V45.1 (old code), V45.11 V45.12, V56.\*; ICD-10: I12.0, I13.11, I13.2, N18.5, N18.6, N19., Z48.22, Z49.\*, Z91.15, Z94.0, Z99.2 [BGP ESRD PMS DXS]
  - SNOMED data set PXRME END STAGE RENAL DISEASE (Problem List only)
- Procedure ICD-9: 38.95, 39.27, 39.42, 39.43, 39.53, 39.93 through 39.95, 54.98, 55.6\*; ICD-10 5A1D70Z, 5A1D80Z, 5A1D90Z [BGP ESRD PROCS]

**Pregnancy Definition**

Any of the following:

- The Currently Pregnant field in Reproductive Factors file set to “Yes” during the Report Period.
- At least one visit during the Report Period, where the primary provider is not a CHR (Provider code 53) with any of the following:
  - Diagnosis (POV or active Problem List entry if added in the past 20 months) ICD-10: O00.1 through O00.91, O09.00 through O10.019, O10.111 through O10.119, O10.211 through O10.219, O10.311 through O10.319, O10.411 through O10.419, O10.911 through O10.919, O11.1 through O15.03, O16.1 through O24.019, O24.111 through O24.119, O24.311 through O24.319, O24.41\*, O24.811 through O24.819, O24.911 through O24.919, O25.10 through O25.13, O26.00 through O26.619, O26.711 through O26.719, O26.811 through O26.93, O28.\*, O29.011 through O30.93, O31.\* through O36.73X9, O36.812 through O48.\*, O60.0\*, O71.00 through O71.03, O88.011 through O88.019, O88.111 through O88.119, O88.211 through O88.219, O88.311 through O88.319, O88.811 through O88.819, O90.3, O91.011 through O91.019, O91.111 through O91.119, O91.211 through O91.219, O92.011 through O92.019, O92.11\*, O92.20, O92.29, O98.011 through O98.019, O98.111 through O98.119, O98.211 through O98.219, O98.311 through O98.319, O98.411 through O98.419, O98.511 through O98.519, O98.611 through O98.619, O98.711 through O98.719, O98.811 through O98.819, O98.911 through

O98.919, O99.011 through O99.019, O99.111 through O99.119, O99.210 through O99.213, O99.280 through O99.283, O99.310 through O99.313, O99.320 through O99.323, O99.330 through O99.333, O99.340 through O99.343, O99.350 through O99.353, O99.411 through O99.419, O99.511 through O99.519, O99.611 through O99.619, O99.711 through O99.719, O99.810, O99.820, O99.830, O99.840 through O99.843, O99.89, O9A.111 through O9A.119, O9A.211 through O9A.219, O9A.311 through O9A.319, O9A.411 through O9A.419, O9A.511 through O9A.519, Z33.1, Z33.3, Z34.\*, Z36.\* [BGP PREGNANCY DIAGNOSES 2]

- CPT 59000-59076, 59300, 59320, 59400-59426, 59510, 59514, 59610, 59612, 59618, 59620, 76801-76828 [BGP PREGNANCY CPT CODES]
- Miscarriage or abortion (see definitions below)

Pharmacy-only visits (clinic code 39) will not count toward this visit. If the patient has more than one pregnancy-related visit during the Report Period, CRS will use the first visit in the Report Period.

- **Miscarriage definition:**

- POV ICD-10: O03.9 [BGP MISCARRIAGE/ABORTION DXS]
- CPT 59812, 59820, 59821, 59830 [BGP CPT MISCARRIAGE]

- **Abortion definition:**

- POV ICD-10: O00.\* through O03.89, O04.\*, Z33.2 [BGP MISCARRIAGE/ABORTION DXS]
- CPT 59100, 59120, 59130, 59136, 59150, 59151, 59840, 59841, 59850, 59851, 59852, 59855, 59856, 59857, S2260 through S2267 [BGP CPT ABORTION]
- Procedure ICD-10: 0WHR73Z, 0WHR7YZ, 10A0\*\*\*, 3E1K78Z, 3E1K88Z [BGP ABORTION PROCEDURES]

## BP Values

Exclusions: When calculating all BPs, the following visits will be excluded:

- Service Category H (Hospitalization), I (In Hospital), S (Day Surgery), or O (Observation)
- Clinic code 23 (Surgical), 30 (ER), 44 (Day Surgery), 79 (Triage), C1 (Neurosurgery), or D4 (Anesthesiology)

CRS uses the last blood pressure documented during the Report Period. If a patient has more than one blood pressure documented on the same day, CRS will first look for a blood pressure less than (<) 140/90 on that day, and if not found, will look for a blood pressure less than (<) 150/90.

### 2.6.2.6 GPRA 2025 Target

During GPRA Year 2025, achieve the target rate of 48.2% for the proportion of patients with blood pressure less than ( $<$ ) 140/90.

### 2.6.2.7 Patient Lists

- List of hypertensive patients with BP less than ( $<$ ) 140/90.
- List of hypertensive patients with BP greater than or equal to ( $\geq$ ) 140/90.

## 2.6.3 Statin Therapy for the Prevention and Treatment of Cardiovascular Disease

### 2.6.3.1 Owner and Contact

Dr. Dena Wilson; Chris Lamer, PharmD

### 2.6.3.2 National Reporting

NATIONAL (included in National GPRA/GPRAMA Report; reported to OMB and Congress)

### 2.6.3.3 Denominators

1. GPRA: User Population patients ages 40 through 75 years with diabetes or a 10-year ASCVD risk score of greater than or equal to ( $\geq$ ) 20% or patients any age with documented ASCVD or patients age 20 through 75 with an LDL greater than or equal to ( $\geq$ ) 190 or an active diagnosis of familial hypercholesterolemia.
2. User Population patients ages 39 years and under with documented ASCVD or ages 20 through 39 with an LDL greater than or equal to ( $\geq$ ) 190 or an active diagnosis of familial hypercholesterolemia.
3. User Population patients ages 40 through 75 years with a 10-year ASCVD risk score of greater than or equal to ( $\geq$ ) 20% or documented ASCVD or an LDL greater than or equal to ( $\geq$ ) 190 or an active diagnosis of familial hypercholesterolemia.
4. User Population patients age 76 years and older with documented ASCVD.
5. User Population patients ages 40 through 75 years with diabetes with LDL less than ( $<$ ) 190, no ASCVD or active diagnosis of familial hypercholesterolemia, and no 10-year ASCVD risk score of greater than or equal to ( $\geq$ ) 20%.

6. User Population patients ages 40 through 75 years with diabetes or a 10-year ASCVD risk score of greater than or equal to ( $\geq$ ) 20% or patients any age with documented ASCVD or patients age 20 through 75 with an LDL greater than or equal to ( $\geq$ ) 190 or an active diagnosis of familial hypercholesterolemia, including denominator exclusions and exceptions.

#### 2.6.3.4 Numerators

1. GPRA: Patients who are statin therapy users during the Report Period or who receive an order (prescription) to receive statin therapy at any point during the Report Period.
2. Patients with any of the listed denominator exclusions or exceptions (with Denominator 6).
  - A. Patients with statin-associated muscle symptoms or an allergy to statin medication.

#### 2.6.3.5 Definitions

##### Diabetes

First DM POV recorded in the V POV file or Problem List Entry with Date of Onset or Date Entered prior to the Report Period:

- ICD-9: 250.00 through 250.93 or ICD-10: E10.\* through E13.\*  
[SURVEILLANCE DIABETES]
- SNOMED data set PXR DIABETES (Problem List only)

##### Atherosclerotic Cardiovascular Disease (ASCVD)

Atherosclerotic Cardiovascular Disease (ASCVD) defined as an active diagnosis of any of the following (POV or Problem List entry where the status is not Inactive or Deleted), or an ASCVD procedure (CABG, PCI, or Carotid Intervention) ever:

- Myocardial Infarction (MI) defined as any of the following:
  - Diagnosis:
    - ICD-9: 410.\*; ICD-10: I21.\* (excluding I21.A1), I22.0, I22.1, I22.8, I22.9 [BGP ECQM MI DXS]
    - SNOMED data set PXR BGP ECQM MI
- Ischemic Heart Disease or Other Related Diagnoses defined as any of the following:
  - Diagnosis:

- ICD-9: 411.\*, 414.2, 414.8, 414.9; ICD-10: I23.0-I23.6, I23.8, I24.0\*, I24.8, I24.9, I25.5, I25.6, I25.82, I25.89, I25.9 [BGP ECQM ISCHEM HEART DIS DXS]
- SNOMED data set PXR M BGP ECQM ISCHEM HEART DIS
- Cerebrovascular Disease, Ischemic Stroke, or Transient Ischemic Attack (TIA) defined as any of the following:
  - Diagnosis:
    - ICD-9: 433.\*, 434.\*, 435.0, 435.1, 435.3, 435.8, 435.9, 436, 437.1, 438.\*, V12.54; ICD-10: G45.0 through G45.2, G45.8, G45.9, G46.0 through G46.8, I63.\*, I69.\*, Z86.73 [BGP ECQM CD STROKE TIA DXS]
    - SNOMED data set PXR M BGP ECQM CD STROKE TIA
- Atherosclerosis and Peripheral Arterial Disease defined as any of the following:
  - Diagnosis:
    - ICD-9: 414.0\*, 414.3, 414.4, 437.0, 440.\*; ICD-10: E08.51, E08.52, E09.51, E09.52, I25.1\*, I25.2, I25.5, I25.700 through I25.812, I25.82 through I25.9, I65.\*, I67.2, I70.\* [BGP ECQM ARTERIAL DISEASE DXS]
    - SNOMED data set PXR M BGP ECQM ARTERIAL DIS
- Stable and Unstable Angina defined as any of the following:
  - Diagnosis:
    - ICD-9: 411.0, 411.1, 413.0, 413.9, 429.79; ICD-10: I20.0, I20.1, I20.8, I20.9, I23.7 [BGP ECQM ANGINA DXS]
    - SNOMED data set PXR M BGP ECQM ANGINA
- Coronary Artery Bypass Graft (CABG) Procedure defined as any of the following:
  - CPT 33510 through 33514, 33516 through 33519, 33521 through 33523, 33533 through 33536, S2205 through S2209
  - Procedure ICD-9: 36.1\*; ICD-10: 02100\*\*, 02103\*\*, 02104\*\*, 02110\*\*, 02113\*\*, 02114\*\*, 02120\*\*, 02123\*\*, 02124\*\*, 02130\*\*, 02133\*\*, 02134\*\* [BGP ECQM CABG PROCS]
  - SNOMED data set PXR M BGP ECQM CABG
- Percutaneous Coronary Interventions (PCI) Procedure defined as any of the following:
  - CPT 92920, 92924, 92928, 92933, 92937, 92941, 92943, 92980 (old code), 92982 (old code), 92995 (old code) [BGP PCI CPTS]

- Procedure ICD-10: 02703\*\*, 02704\*\*, 02713\*\*, 02714\*\*, 02723\*\*, 02724\*\*, 02733\*\*, 02734\*\* [BGP ECQM PCI PROCS]
- SNOMED data set PXRMBGP ECQM PCI
- Carotid Intervention defined as any of the following:
  - Procedure ICD-9: 00.61 through 00.65, 38.02, 38.12, 38.22, 38.30 through 38.32, 38.42, 39.22, 39.28, 88.41; ICD-10: 03\*H\*\*\*, 03\*J\*\*\*, 03\*K\*\*\*, 03\*L\*\*\*, 03\*M\*\*\*, 03\*N\*\*\*, 0G\*6\*\*\*, 0G\*7\*\*\*, 0G\*8\*\*\*, B3060ZZ, B3061ZZ, B306YZZ, B3070ZZ, B3071ZZ, B307YZZ, B3080ZZ, B3081ZZ, B308YZZ, B3160ZZ, B3161ZZ, B316YZZ, B3170ZZ, B3171ZZ, B317YZZ, B3180ZZ, B3181ZZ, B318YZZ [BGP ECQM CAROTID INTER PROCS]
  - SNOMED data set PXRMBGP ECQM CAROTID INTER

## LDL

For LDL greater than or equal to ( $\geq$ ) 190, CRS will look for any test at any time with a result greater than or equal to ( $\geq$ ) 190. LDL is defined as any of the following:

- LOINC taxonomy: 12773-8, 13457-7, 18261-8, 18262-6, 2089-1, 2090-9, 22748-8, 35198-1, 39469-2, 49132-4, 55440-2, 69419-0, 96258-9, 96259-7, 96597-0 [BGP LDL LOINC CODES]
- Site-populated taxonomy DM AUDIT LDL CHOLESTEROL TAX

## Familial hypercholesterolemia

Diagnosis (POV or Problem List entry where the status is not Inactive or Deleted):

- POV ICD-10: E78.01 [BGP ECQM HYPERCHOL DXS]
- SNOMED data set PXRMBGP ECQM HYPERCHOL

## 10-year ASCVD risk score

Any of the following during the report period:

- Measurement ACC (10 YEAR ASCVD RISK)

## Denominator Exclusions

Patients meeting any of the following conditions will be excluded from the denominator:

- Patients who are breastfeeding.
- Patients who have a diagnosis of rhabdomyolysis during the report period (POV or Problem List entry where the status is not Inactive or Deleted).

## Denominator Exceptions

The patient is excluded from the denominator if they meet any of the following criteria and do not meet the numerator.

- Patients with statin-associated muscle symptoms or an allergy to statin medication during the Report Period.
- Patients with active liver disease or hepatic disease (including Hepatitis A or B [POV or Problem List entry where the status is not Inactive or Deleted]) or insufficiency.
- Patients who are receiving hospice or palliative care during the Report Period.
- Patients with end-stage renal disease (ESRD).
- Patients with documentation of a medical reason for not being prescribed statin therapy during the Report Period.

## Statin-associated Muscle Symptoms or an Allergy

Defined as any of the following during the Report Period:

- POV ICD-9: 359.9, 729.1; ICD-10: G72.0, G72.9, M60.9, M79.10
- SNOMED data set PXRMBGP ECQM STATIN ADV
- POV ICD-9: 995.0 through 995.3 [BGP ASA ALLERGY 995.0-995.3] and E942.9 [BGP ADV EFF CARDIOVASC NEC]
- “Statin” or “Statins” entry (except “Nystatin”) in ART (Patient Allergies File)
- “Statin” or “Statins” (except “Nystatin”) contained within Problem List (where status is not Deleted) or in Provider Narrative field for any POV ICD-9: 995.0 through 995.3, V14.8; ICD-10: Z88.8 [BGP ASA ALLERGY 995.0-995.3 and BGP HX DRUG ALLERGY NEC]
- Problem List entry where the status is not Deleted of SNOMED data set PXRMBGP ADR STATIN
- NMI (not medically indicated) refusal for any statin at least once during the Report Period

## Breastfeeding Definition

Any of the following documented during the Report Period:

- The Lactation Status field in Reproductive Factors file set to “Lactating”
- Diagnosis:
  - POV ICD-10: O91.03, O91.13, O91.23, O92.03, O92.13, O92.5, O92.70, O92.79, Z39.1 [BGP ECQM BREASTFEED DXS]
  - SNOMED data set PXRMBGP ECQM BREASTFEED

- Breastfeeding Patient Education codes BF-BC, BF-BP, BF-CS, BF-EQ, BF-FU, BF-HC, BF-ON, BF-M, BF-MK, BF-N or containing SNOMED data sets PXRMBGPPTEDBREASTFEED or PXRMBGP ECQM BREASTFEED

### **Rhabdomyolysis**

- Diagnosis (POV or Problem List entry where the status is not Inactive or Deleted):
  - ICD-10: M62.82, T79.6XX\* [BGP ECQM RHABDO DXS]
  - SNOMED data set PXRMBGP ECQM RHABDO (Problem List only)

### **Active Liver Disease, Hepatic Disease or Insufficiency Definition**

Diagnosis (POV or Problem List entry where the status is not Deleted):

- POV ICD-10: B17.\*, B18.2 through B19.0, B19.20, B19.21, B19.9, K70.0 through K74.69, K75.4, O98.41\* [BGP ECQM LIVER DISEASE DXS]
- SNOMED data set PXRMBGP ECQM LIVER DIS

### **Hospice Indicator**

Hospice is defined as any of the following during the Report Period:

- SNOMED data set PXRMBGP IPC INPT ENC (Inpatient encounter) with DISCHARGE SNOMED CT data set PXRMBGP IPC DISCHG HOSPICE (discharge to home or health care facility for hospice care).
- Visit, V CPT, V POV or Problem List entry where the status is not Inactive or Deleted: SNOMED data set PXRMBGP IPC HOSPICE (hospice care ambulatory, hospice encounter, hospice diagnosis).
- CPT 99377, 99378, G0182, G9473, G9474, G9475, G9476, G9477, G9478, G9479, Q5003, Q5004, Q5005, Q5006, Q5007, Q5008, Q5010, S9126, T2042, T2043, T2044, T2045, T2046 [BGP ECQM HOSPICE CPTS].

### **Palliative Care**

Palliative care is defined as any of the following during the Report Period:

- CPT G9054
- SNOMED data set PXRMBGP ECQM PALLIATIVE ENC
- POV ICD-10: Z51.5



**ESRD**

End Stage Renal Disease diagnosis or treatment is defined as any of the following ever:

- CPT 36145 (old code), 36147, 36800, 36810, 36815, 36818, 36819, 36820, 36821, 36831 through 36833, 50300, 50320, 50340, 50360, 50365, 50370, 50380, 90951 through 90970 or old codes 90918 through 90925, 90935, 90937, 90939 (old code), 90940, 90945, 90947, 90989, 90993, 90997, 90999, 99512, 3066F, G0257, G9231, M1187, M1188, S2065, S9339 [BGP ESRD CPTS]
- Diagnosis (POV or Problem List entry where the status is not Inactive or Deleted):
  - ICD-9: 585.6, V42.0, V45.1 (old code), V45.11, V45.12, V56.\*; ICD-10: I12.0, I13.11, I13.2, N18.5, N18.6, N19., Z48.22, Z49.\*, Z91.15, Z94.0, Z99.2 [BGP ESRD PMS DXS]
  - SNOMED data set PXRMM END STAGE RENAL DISEASE (Problem List only)
- Procedure ICD-9: 38.95, 39.27, 39.42, 39.43, 39.53, 39.93 through 39.95, 54.98, 55.6\*; ICD-10 5A1D70Z, 5A1D80Z, 5A1D90Z [BGP ESRD PROCS]

**Hepatitis A Definition**

Diagnosis (POV or Problem List entry where the status is not Deleted):

- ICD-10: B15.\* [BGP IPC HEP A DXS]
- SNOMED data set PXRMM BGP IPC HEP A EVID

**Hepatitis B Definition**

Diagnosis (POV or Problem List entry where the status is not Deleted):

- ICD-10: B16.\*, B18.0, B18.1, B19.1\*[BGP ECQM HEP B DXS]
- SNOMED data set PXRMM BGP ECQM HEP B

**Medical Reason for Not Being Prescribed Statin Therapy**

- SNOMED data set PXRMM BGP IPC NOT DONE MED for statin medication not ordered.

**Statins Numerator Logic**

- **Statin Therapy Users**
  - CPT 4013F
- **Statin Medication Codes**
  - Defined with medication taxonomy BGP PQA STATIN MEDS.

- **Statin medications and combination products are:** atorvastatin (+/- amlodipine, ezetimibe), fluvastatin, lovastatin (+/- niacin), pitavastatin, pravastatin, rosuvastatin (+/- ezetimibe), simvastatin (+/- ezetimibe, niacin, sitagliptin).

Patients must have an active prescription for statin therapy during the Report Period. This includes patients who receive an order during the Report Period, or prior to the Report Period with enough days' supply to take them into the Report Period.

$$\text{Rx Days' Supply} \geq (\text{Report Period Begin Date} - \text{Prescription Date})$$

Active prescriptions include active outside medications, defined as V Med entry at any time with EHR OUTSIDE MED field not blank and DATE DISCONTINUED field blank.

### 2.6.3.6 GPRA 2025 Target

During GPRA Year 2025, achieve the target rate of 36.9% for the proportion of at-risk patients who receive statin therapy.

### 2.6.3.7 Patient Lists

- List of patients 40–75 years old with diabetes or any age with ASCVD or 20-75 years old with LDL greater than or equal to ( $\geq$ ) 190 or familial hypercholesterolemia or 40-75 years old with a 10-year ASCVD risk score greater than or equal to ( $\geq$ ) 20% with statin therapy.
- List of patients 40–75 years old with diabetes or any age with ASCVD or 20-75 years old with LDL greater than or equal to ( $\geq$ ) 190 or familial hypercholesterolemia or 40-75 years old with a 10-year ASCVD risk score greater than or equal to ( $\geq$ ) 20% without statin therapy.
- List of patients with denominator exclusions.

## 2.7 STD-Related Group

### 2.7.1 HIV Screening

#### 2.7.1.1 Owner and Contact

Richard Haverkate, MPH

#### 2.7.1.2 National Reporting

NATIONAL (included in IHS Performance Report; reported to OMB and Congress)

### 2.7.1.3 Denominators

1. Pregnant Active Clinical patients with no documented miscarriage or abortion during the past 20 months and *no* recorded HIV diagnosis ever.
2. GPRA: User Population patients ages 13–64 years with no recorded HIV diagnosis prior to the Report Period.

### 2.7.1.4 Numerators

**Note:** These numerators do *not* include refusals.

1. Patients who were screened for HIV during the past 20 months (with Denominator 1).
2. GPRA: Patients who were screened for HIV at any time before the end of the Report Period (with Denominator 2).

### 2.7.1.5 Definitions

#### HIV

Any of the following documented any time prior to the end of the Report Period:

- POV or Problem List entry where the status is not Deleted:
  - ICD-9: 042, 079.53, V08; ICD-10: B20, B97.35, Z21, O98.711 through O98.73 [BGP HIV/AIDS DXS]
  - SNOMED data set PXRMI HIV (Problem List only)

#### Pregnancy

Any of the following:

- The Currently Pregnant field in Reproductive Factors file set to “Yes” during the Report Period.
- At least two visits during the past 20 months from the end of the Report Period, where the primary provider is not a Community Health Representative (CHR) (Provider code 53) with any of the following:

- Diagnosis (POV or active Problem List entry if added in the past 20 months) ICD-10: O00.1 through O00.91, O09.00 through O10.019, O10.111 through O10.119, O10.211 through O10.219, O10.311 through O10.319, O10.411 through O10.419, O10.911 through O10.919, O11.1 through O15.03, O16.1 through O24.019, O24.111 through O24.119, O24.311 through O24.319, O24.41\*, O24.811 through O24.819, O24.911 through O24.919, O25.10 through O25.13, O26.00 through O26.619, O26.711 through O26.719, O26.811 through O26.93, O28.\*, O29.011 through O30.93, O31.\* through O36.73X9, O36.812 through O48.\*, O60.0\*, O71.00 through O71.03, O88.011 through O88.019, O88.111 through O88.119, O88.211 through O88.219, O88.311 through O88.319, O88.811 through O88.819, O90.3, O91.011 through O91.019, O91.111 through O91.119, O91.211 through O91.219, O92.011 through O92.019, O92.11\*, O92.20, O92.29, O98.011 through O98.019, O98.111 through O98.119, O98.211 through O98.219, O98.311 through O98.319, O98.411 through O98.419, O98.511 through O98.519, O98.611 through O98.619, O98.711 through O98.719, O98.811 through O98.819, O98.911 through O98.919, O99.011 through O99.019, O99.111 through O99.119, O99.210 through O99.213, O99.280 through O99.283, O99.310 through O99.313, O99.320 through O99.323, O99.330 through O99.333, O99.340 through O99.343, O99.350 through O99.353, O99.411 through O99.419, O99.511 through O99.519, O99.611 through O99.619, O99.711 through O99.719, O99.810, O99.820, O99.830, O99.840 through O99.843, O99.89, O9A.111 through O9A.119, O9A.211 through O9A.219, O9A.311 through O9A.319, O9A.411 through O9A.419, O9A.511 through O9A.519, Z33.1, Z33.3, Z34.\*, Z36.\* [BGP PREGNANCY DIAGNOSES 2]
- CPT 59000-59076, 59300, 59320, 59400-59426, 59510, 59514, 59610, 59612, 59618, 59620, 76801-76828 [BGP PREGNANCY CPT CODES]

Pharmacy-only visits (clinic code 39) will not count toward these two visits. If the patient has more than two pregnancy-related visits during the past 20 months, CRS will use the first two visits in the 20-month period. In addition, the patient must have at least one pregnancy-related visit occurring during the reporting period.

The patient must not have a documented miscarriage or abortion occurring after the second pregnancy-related visit or the date the Currently Pregnant field was set to “Yes.” The period is extended to include patients who were pregnant during the Report Period, but whose initial diagnosis (and HIV test) were documented prior to Report Period.

- **Miscarriage definition**

- POV ICD-10: O03.9 [BGP MISCARRIAGE/ABORTION DXS]

- CPT 59812, 59820, 59821, 59830 [BGP CPT MISCARRIAGE]
- **Abortion definition**
  - POV ICD-10: O00.\* through O03.89, O04.\*, Z33.2 [BGP MISCARRIAGE/ABORTION DXS]
  - CPT 59100, 59120, 59130, 59136, 59150, 59151, 59840, 59841, 59850, 59851, 59852, 59855, 59856, 59857, S2260 through S2267 [BGP CPT ABORTION]
  - Procedure ICD-10: 0WHR73Z, 0WHR7YZ, 10A0\*\*\*, 3E1K78Z, 3E1K88Z [BGP ABORTION PROCEDURES]

### HIV Screening

- CPT 80081, 86689, 86701 through 86703, 87389 through 87391, 87534 through 87539, 87806, 87901, 87906 [BGP CPT HIV TESTS]
- LOINC taxonomy: 10901-7, 10902-5, 11078-3, 11079-1, 11080-9, 11081-7, 11082-5, 12855-3, 12856-1, 12857-9, 12858-7, 12859-5, 12870-2, 12871-0, 12872-8, 12875-1, 12876-9, 12893-4, 12894-2, 12895-9, 13499-9, 13920-4, 14092-1, 14126-7, 16132-3, 16974-8, 16975-5, 16978-9, 18396-2, 19110-6, 21007-0, 21009-6, 21331-4, 21332-2, 21334-8, 21335-5, 21336-3, 21337-1, 21338-9, 21339-7, 21340-5, 22356-0, 22357-8, 22358-6, 23876-6, 24012-7, 28004-0, 28052-9, 29327-4, 29893-5, 30245-5, 30361-0, 31072-2, 31073-0, 31201-7, 31430-2, 32571-2, 32602-5, 32827-8, 32842-7, 33508-3, 33660-2, 33806-1, 33807-9, 33866-5, 34591-8, 34592-6, 34699-9, 35437-3, 35438-1, 35439-9, 35440-7, 35441-5, 35442-3, 35443-1, 35444-9, 35445-6, 35446-4, 35447-2, 35448-0, 35449-8, 35450-6, 35452-2, 35564-4, 35565-1, 38998-1, 40437-6, 40438-4, 40439-2, 40732-0, 40733-8, 41143-9, 41144-7, 41145-4, 41290-8, 42339-2, 42600-7, 42627-0, 42768-2, 43008-2, 43009-0, 43010-8, 43011-6, 43012-4, 43013-2, 43185-8, 43599-0, 44531-2, 44532-0, 44533-8, 44607-0, 44871-2, 44872-0, 44873-8, 45212-8, 47029-4, 48023-6, 48345-3, 48346-1, 49483-1, 49580-4, 49718-0, 49905-3, 49965-7, 51786-2, 51866-2, 5220-9, 5221-7, 5222-5, 5223-3, 5224-1, 5225-8, 53379-4, 53601-1, 53825-6, 54086-4, 56888-1, 57974-8, 57975-5, 57976-3, 57977-1, 57978-9, 58900-2, 59052-1, 59419-2, 62456-9, 6429-5, 6430-3, 6431-1, 68961-2, 69668-2, 73905-2, 73906-0, 74856-6, 75622-1, 75666-8, 77685-6, 7917-8, 7918-6, 7919-4, 80203-3, 80387-4, 81641-3, 83101-6, 85037-0, 85361-4, 85368-9, 85686-4, 86233-4, 88453-6, 89365-1, 89374-3, 95524-5, 95523-7, 96273-8, 96556-6, 96557-4, 9660-2, 9661-0, 9662-8, 9663-6, 9664-4, 9665-1, 9666-9, 9667-7, 9668-5, 9669-3, 97860-1, 97861-9, 9821-0, 9836-8, 9837-6, 105080-6 [BGP HIV TEST LOINC CODES]
- Site-populated taxonomy BGP HIV TESTS

**Note:** The time frame for screening for the pregnant patient's denominator is anytime during the past 20 months.

### 2.7.1.6 GPRA 2025 Target

During GPRA Year 2025, achieve the target rate of 42.5% for the proportion of patients who have ever been screened for HIV.

### 2.7.1.7 Patient Lists

- List of pregnant patients with documented HIV test in the past 20 months.
- List of pregnant patients without documented HIV test in the past 20 months.
- List of User Population patients 13–64 years old with documented HIV test ever.
- List of User Population patients 13–64 years old without documented HIV test ever.

## 2.8 Other Clinical Group

### 2.8.1 Breastfeeding Rates

**Note:** This measure is used in conjunction with the Childhood Weight Control GPRA long-term measure to support the reduction of the incidence of childhood obesity.

#### 2.8.1.1 Owner and Contact

Tina Tah, RN, BSN, MBA

#### 2.8.1.2 National Reporting

NATIONAL (included in IHS Performance Report; reported to OMB and Congress)

#### 2.8.1.3 Denominators

1. Active Clinical patients who are 30 through 394 days old who were screened for infant feeding choice at the age of 2 months (38 through 89 days).
2. User Population patients ages 30 through 394 days old.
3. GPRA: User Population patients who are 30 through 394 days old who were screened for infant feeding choice at the age of 2 months (38 through 89 days).
4. User Population patients 30 through 394 days old who were screened for infant feeding choice at the age of 6 months (165 through 209 days).
5. User Population patients 30 through 394 days old who were screened for infant feeding choice at the age of 9 months (255 through 299 days).

6. User Population patients 30 through 394 days old who were screened for infant feeding choice at the age of 1 year (350 through 394 days).

#### **2.8.1.4 Numerators**

1. Patients who were screened for infant feeding choice at least once (with Denominator 1).
2. Patients who were screened for infant feeding choice at the age of 2 months (38 through 89 days) (with Denominator 1).
3. Patients who were screened for infant feeding choice at the age of 6 months (165 through 209 days) (with Denominator 1).
4. Patients who were screened for infant feeding choice at the age of 9 months (255 through 299 days) (with Denominator 1).
5. Patients who were screened for infant feeding choice at the age of 1 year (350 through 394 days) (with Denominator 1).
6. GPRA: Patients who, at the age of 2 months (38 through 89 days), were either exclusively or mostly breastfed (with Denominator 2).
7. Patients who, at the age of 6 months (165 through 209 days), were either exclusively or mostly breastfed (with Denominator 3).
8. Patients who, at the age of 9 months (255 through 299 days), were either exclusively or mostly breastfed (with Denominator 4).
9. Patients who, at the age of 1 year (350 through 394 days), were either exclusively or mostly breastfed (with Denominator 5).

#### **2.8.1.5 Definitions**

##### **Patient Age**

Since the age of the patient is calculated at the beginning of the Report Period, this measure may include patients up to 25 months old if they were within the eligible age range on the first day of the Report Period. It will *not* include any patients who were born after the first day of the Report Period. Patients born after the first day of the Report Period will be included in the following Report Period.

## Infant Feeding Choice

The documented feeding choice from the V Infant Feeding Choice file that is closest to the exact age that is being assessed will be used. For example, if a patient was assessed at 45 days old as half breastfed and half formula, then assessed again at 65 days old as mostly breastfed, the mostly breastfed value will be used since it is closer to the exact age of 2 months (i.e., 60 days). Another example is a patient who was assessed at 67 days as mostly breastfed and again at 80 days as mostly formula. In this case, the 67-day value of mostly breastfed will be used. The other exact ages are 180 days for 6 months, 270 days for 9 months, and 365 days for one year.

To be included in the age-specific screening numerators, the patient must have been screened within the specific age range. For example, if a patient was screened at 6 months and was exclusively breastfed, but was not screened at 2 months, then the patient will only be counted in the 6 months numerator.

### 2.8.1.6 GPRA 2025 Target

During GPRA Year 2025, achieve the target rate of 44.1% for the proportion of two-month-olds who are mostly or exclusively breastfeeding.

### 2.8.1.7 Patient Lists

- List of patients ages 30 through 394 days who were screened for Infant Feeding Choice at least once.
- List of patients ages 30 through 394 days who were not screened for Infant Feeding Choice at least once.
- List of User Population patients screened at the age of 2 months (38 through 89 days) and were either exclusively or mostly breastfed.
- List of User Population patients screened at the age of 2 months (ages 38 through 89 days) and were not exclusively or mostly breastfed.



## Acronyms

| Acronym  | Term Meaning   |
|----------|--|
| ADA      | American Diabetes Association  |
| AI/AN    | American Indian/Alaska Native  |
| AMI      | Acute Myocardial Infarction  |
| BH       | Behavioral Health  |
| BHS      | Behavioral Health System   |
| BI       | Brief Intervention   |
| BMI      | Body Mass Index  |
| BNI      | Brief Negotiated Interview   |
| BP       | Blood Pressure   |
| CABG     | Coronary Artery Bypass Graft   |
| CAGE     | Cut Down, Annoyed, Guilty, and Eye Opener (alcohol use disorders screening test)   |
| CAGE-AID | CAGE-Adapted to Include Drugs (alcohol and substance use disorders screening test) |
| CHR      | Community Health Representative  |
| CPT      | Current Procedural Terminology   |
| CRC      | Colorectal Cancer  |
| CRS      | Clinical Reporting System  |
| CVD      | Cardiovascular Disease   |
| CVX      | Vaccine Code   |
| DBH      | Division of Behavioral Health  |
| DCBE     | Double Contrast Barium Enema   |
| DM       | Diabetes Mellitus  |
| DNKA     | Did Not Keep Appointment   |
| DTaP     | Diphtheria Tetanus Acellular Pertussis   |
| DTP      | Diphtheria, Tetanus and Pertussis  |
| ER       | Emergency Room   |
| ESRD     | End Stage Renal Disease  |
| ETDRS    | Early Treatment Diabetic Retinopathy Study   |
| ETS      | Environmental Tobacco Smoke  |
| FIT      | Fecal Immunochemical Test  |
| FOBT     | Fecal Occult Blood Test  |
| FY       | Fiscal Year  |
| GFR      | Glomerular Filtration Rate   |
| GPRA     | Government Performance and Results Act of 1993                                     |

| <b>Acronym</b> | <b>Term Meaning</b>                                      |
|----------------|--|
| GPRAMA         | GPRA Modernization Act                                   |
| HEDIS          | Healthcare Effectiveness Data and Information Set        |
| HGB            | Hemoglobin   |
| HHS            | Health and Human Services                                |
| HiB            | Haemophilus Influenza Type B                             |
| HIV            | Human Immunodeficiency Virus                             |
| HPV            | Human Papillomavirus                                     |
| ICD            | International Classification of Diseases                 |
| IFC            | Infant Feeding Choice                                    |
| IHS            | Indian Health Service                                    |
| IMM            | Immunization   |
| IPV            | Inactivated Polio Vaccine                                |
| IPV/DV         | Intimate Partner Violence/Domestic Violence              |
| LCSW           | Licensed Clinical Social Worker                          |
| LDL            | Low-Density Lipoprotein                                  |
| LOINC          | Logical Observations Identifiers, Names, Codes           |
| MMR            | Measles, Mumps, Rubella (Vaccine)                        |
| MPH            | Master's in Public Health                                |
| M/R            | Measles and Rubella                                      |
| NHANES         | National Health and Nutrition Examination Survey         |
| NMI            | Not Medically Indicated                                  |
| NOS            | Not Otherwise Specified                                  |
| NQF            | National Quality Forum                                   |
| OMB            | Office of Management and Budget                          |
| OPV            | Oral Polio Vaccine                                       |
| PCC            | Patient Care Component                                   |
| PCI            | Percutaneous Coronary Interventions                      |
| POV            | Purpose of Visit   |
| PRC            | Purchased and Referred Care                              |
| R/M            | Rubella/Mumps  |
| RPMS           | Resource and Patient Management System                   |
| SBIRT          | Screening, Brief Intervention, and Referral to Treatment |
| SNOMED         | Systematized Nomenclature of Medicine                    |
| STD            | Sexually Transmitted Disease                             |
| Tdap           | Tetanus, Diphtheria and Acellular Pertussis              |
| TIA            | Transient Ischemic Attack                                |
| TUHF           | Tobacco User Health Factors                              |

---

| Acronym | Term Meaning                      |
|---------|-----------------------------------|
| UACR    | Urine Albumin-to-Creatinine Ratio |

## Contact Information

If you have any questions or comments regarding this distribution, please contact the IHS IT Service Desk.

**Phone:** (888) 830-7280 (toll free)

**Web:** <https://www.ihs.gov/itsupport/>

**Email:** [itsupport@ihs.gov](mailto:itsupport@ihs.gov)