

**Exhibit 3-41 I
Annual Competency for Admixing IVs for Personnel Performing Immediate Use
Compounding**

Please Circle **True or False**

True or False	IV medications should only be compounded in the pharmacy (under a laminar flow hood) when there is an emergency or when the product stability is short.
True or False	IV medications made outside of the pharmacy or laminar flow hood should be prepared for immediate-use only, meaning they need to be hung within 4 hours from the start of making the preparation or else disposed of properly. There is no requirement for the duration of administration.
True or False	Before making the IV for immediate use, hand washing or hand rub with sanitizer for hand hygiene should be performed per CDC guidelines.
True or False	In an emergency, it is acceptable to make the Immediate Use IV in a crash cart.
True or False	The designated area for IV medication preparation should include an area that is clean and clutter-free and is the dedicated medication preparation tray.
True or False	The dedicated medication preparation area should be cleaned with bleach, alcohol, or solvent cleaner before each use.
True or False	Any questionable product after inspecting it for particulate matter, cracks, leaks, clarity, color, etc., should be disposed of properly and the process restarted.
True or False	The IV made for immediate use shall include no more than 3 sterile commercially manufactured, non-hazardous products.
True or False	The person that requested you to make the IVs is responsible for labeling the IV.
True or False	High Alert medications used to prepare an Immediate Use IV require a 2nd check verification completed per policy.

Please list what is required on the Immediate Use IV label:

Signature of Person Assessed: _____ Date: _____

Printed Name: _____

Signature of Qualified Evaluator: _____ Date: _____

Printed Name: _____