

Exhibit 3-41 L

Frequency for Minimum Cleaning and Disinfection of Classified Areas and in the Segregated Compounding Area

Site	Cleaning	Disinfecting	Applying Sporocidal	Example
ISO 5 Class PEC and equipment inside PEC	Equipment and all interior surfaces of the PEC daily on days when compounding occurs and when surface contamination is known or suspected	Equipment and all interior surfaces of the PEC daily on days when compounding occurs and when surface contamination is known or suspected	Monthly	Quaternary Ammonium, 70% Sterile IPA, hydrogen peroxide/peracetic acid (sporicidal)
Removable work tray of the PEC, when applicable	-Work surface of the tray daily on days when compounding occurs -All surfaces of the area underneath the work tray monthly	-Work surface of the tray daily on days when compounding occurs -All surfaces of the area underneath the work tray monthly	Monthly for all surfaces	
Pass-through chambers	Daily on days when compounding occurs	Daily on days when compounding occurs	Monthly	
Work surface(s) outside the PEC	Daily on days when compounding occurs	Daily on days when compounding occurs		
Floor(s)	Daily on days when compounding occurs	Daily on days when compounding occurs		
Sink(s)	Daily each day in use	Daily each day in use		
Walls, doors, door frames	Monthly	Monthly	Monthly	
Ceilings*	Monthly	Monthly	Monthly	
Storage shelving and bins	Monthly	Monthly	Monthly	
Equipment outside the PECs	Monthly	Monthly	Monthly	

* Ceilings of an SCA are required to be cleaned, disinfected, and applied with sporicidal only when visibly soiled or surface contamination is known or suspected.

+ Consider disinfectants that are registered by the EPA as a one-step disinfectant. This means that the disinfectant has been formulated to be effective in the presence of light to moderate soiling with the pre-cleaning step.

++ Cleaning, disinfecting, and sporicidal agents used in the PEC must be sterile.